arXiv:0712.2533v1 [math.AT] 17 Dec 2007

The Viterbo Transfer as a Map of Spectra

Thomas Kragh[1]

ABSTRACT. Let L and N be two smooth manifolds of the same dimension. Let $j\colon L \to T^*N$ be an exact Lagrange embedding. We denote the free loop space of X by ΛX. In [10], C. Viterbo constructed a transfer map $(\Lambda j)^!\colon H^*(\Lambda L) \to H^*(\Lambda N)$. This transfer is constructed using finite reduction of Floer homology. In this paper we define a continuous family of finite reductions that makes it possible to realize this transfer as a map of Thom spectra: $(\Lambda j)_!\colon (\Lambda N)^{-TN} \to (\Lambda L)^{-TL+\eta}$, where η is a virtual bundle classified by the tangential information of j.

Contents

1 Introduction and Statement of Results . . . 1

2 The Homotopy Index . . . 5

3 The Action Integral in Cotangent Bundles . . . 8

4 Finite Reductions of the Action Integral in Cotangent Bundles for a Hamiltonian Linear at Infinity . . . 9

5 Generalized Finite Reductions . . . 21

6 Stabilization of Generalized Finite Reductions . . . 29

7 The Maslov Bundle and Index of Finite Reductions . . . 34

8 The Viterbo Transfer as a Map of Spectra . . . 42

Bibliography . . . 48

1 Introduction and Statement of Results

Let L and N be closed, n-dimensional, smooth manifolds, and let $\pi\colon T^*N \to N$ be the projection. The Liouville form (or canonical 1-form) λ is defined by

$$\lambda_{q,p}(v) = p(\pi_*(v)), \quad q \in N, p \in T^*_qN, v \in T_{q,p}(T^*N).$$

---

[1] Partially funded by CTQM, OP-ALG-TOP-GEO and The University of Aarhus.

The 2-form $\omega = d\lambda$ is non-degenerate and thus defines a canonical symplectic structure on T^*N. An embedding $j\colon L \to T^*N$ is called Lagrange if $j^*\omega = dj^*\lambda = 0$ and exact Lagrange if $j^*\lambda$ is exact. We assume from now on that j is an exact Lagrange embedding. The trivial examples of such embeddings are those which are Hamiltonian isotopic to the zero section. To this day no non-trivial examples have been found, and the nearby Lagrangian conjecture states that there are no other.

Resently there has been much progress in this area. Specifically in [9] and [5] it is proven, independently and in different ways, that under certain conditions j is a homology equivalence. However, there is still the general case, and there is a big step from being a Homology equivalence to being Hamiltonian isotopic to the zero section.

We denote the free loop space of a space X by ΛX. In [10], Viterbo constructs a transfer map $(\Lambda j)^!$ on cohomology, such that

$$\begin{tikzcd}
H^*(\Lambda L) \arrow[r, "(\Lambda j)^!"] \arrow[d, shift left, "i^*"] & H^*(\Lambda N) \arrow[d, shift left, "i^*"] \\
H^*(L) \arrow[r, "(\pi\circ j)^!"] \arrow[u, shift left, "\mathrm{Ev}_0^*"] & H^*(M) \arrow[u, shift left, "\mathrm{Ev}_0^*"]
\end{tikzcd}$$

commutes. Here $(\pi\circ j)^!$ is the standard transfer map on cohomology, Ev_0 is the evaluation at base point, and i is the inclusion of constant curves. In this paper we call this map the Viterbo transfer. Viterbo uses this transfer as obstruction to the existence of exact Lagrange embeddings.

Because $j\colon L \to T^*N$ is Lagrange we get a Maslov class in $H^1(L)$. This defines a map $\Lambda L \to \mathbb{Z}$ called the Maslov index, and it turns out that the Viterbo transfer is graded on each component by this Maslov index. We prove the following theorem in this paper.

Theorem 1. *The Viterbo transfer can be realized as a map of spectra*

$$(\Lambda j)_!\colon (\Lambda N)^{-TN} \to (\Lambda L)^{-TL+\eta},$$

where η is a virtual bundle classified by the tangential information of the embedding $j\colon L \to T^*N$, and the local dimension of η is the Maslov index.

In the original construction by Viterbo, the Thom isomorphism is used on what turns out to be the virtual bundle $-TL+\eta$. However, η is not necessarily oriented, but if we assume $(\pi\circ j)\colon L \to N$ to be relative spin, it will be. This is part of the conditions in [9] and [5]. In fact, they assume that both N and L are spin and that the Maslov class vanishes. This implies that the virtual bundle η is zero dimensional and oriented. In fact, if we also assume that L and N are simply connected, one can prove that the Viterbo transfer is a homotopy equivalence of spectra, putting restrictions on η. This may yield new insight into the situation, but for now we only construct the transfer.

The rest of this section is devoted to giving an overview of the construction using finite reductions of Floer homology.

Outline of proof of Theorem 1: The actual construction does not use Floer homology. It is, however, very illuminating to sketch the relation. This relation also justifies considering the spectra constructed as representations of the stable homotopy type of Floer homology in cotangent bundles.

For any Hamiltonian $H\colon T^*N \to \mathbb{R}$ we define the action integral

$$A_H\colon \Lambda T^*N \to \mathbb{R}$$

by the formula

$$A_H(\gamma) = \int_\gamma \lambda - H dt.$$

In this paper we will always have $H(q,p) = \mu\|p\|$ for large $\|p\|$, where μ is not a geodesic length. Floer homology $FH_*(T^*N, H)$ is essentially Morse homology of A_H perturbed on the infinite dimensional manifold ΛT^*N (see e.g. [1]). The linear at infinity case can be calculated to be

$$FH_*(T^*N, H) \approx H_*(\Lambda^\mu N), \tag{1}$$

where $\Lambda^\mu N$ is the space of curves with length less than μ. Define $T^*\Lambda_r N$ as the cotangent space of the manifold of r-pieced geodesics each of length less than some ε. The finite reductions we define in this paper can be constructed in the following way: for large r we define embeddings:

$$i_r\colon T^*\Lambda_r N \to \Lambda T^*N,$$

satisfying:

- The image of i_r contains all critical points of A_H
- The composition $A_r = A_H \circ i_r$ has no other critical points than those from A_H
- There is a "consistent" way of doing Morse theory for A_r on $T^*\Lambda_r N$ and creating a space Z such that the homotopy type of this space does not change under small compact pertubations of H.

In the actual construction, we use the theory of homotopy indices described in section 2, but for the purpose of this overview, one may think of A_r as a Morse function, and thus think of Z as a cell complex with one cell per critical point.

In section 4 we explicitly define a function A_r as above and prove that Z is homotopy equivalent to the Thom space

$$\mathrm{Th}(T\Lambda_r^\mu N) = (\Lambda_r^\mu N)^{T\Lambda_r^\mu N} = D(T\Lambda_r^\mu N)/S(T\Lambda_r^\mu N),$$

where $\Lambda_r^\mu N$ is the manifold of piecewise geodesics, with r pieces each having length less than μ/r, and where $D(\cdot)$ is the disc bundle, and $S(\cdot)$ the sphere bundle. This was already proven by Viterbo in [10], but because we have an explicitly defined A_r, we can prove it more directly. Using the Thom isomorphism, this is consistent with equation (1).

Because $j\colon L \to T^*N$ is a Lagrange embedding, we can use the Darboux-Weinstein theorem to extend j to a symplectic embedding of a small neighborhood of the zero section in T^*L. Using this neighborhood and the fact that j is exact, we can adjust H such that all the critical points of A_r with critical values above c, for some c, are curves inside the neighborhood of L. In fact, in the small neighborhood of L minus an even smaller neighborhood of L H

will be defined to be linear with some slope μ_L. We now define Z_L to be the complex defined by the quotient of Z by the subcomplex Y defined by those cells associated with critical points with critical value less than c.

If we were to go through the above construction for H restricted to the neighborhood of L using the cotangent bundle structure of T^*L, we would get a space Z'_L which would be homotopy equivalent to $\mathrm{Th}(T\Lambda_r^{\mu_L}L)$. However, the construction of A_r highly depends on the vertical Lagrangian foliation in T^*N, and in the small neighborhood of L the two foliations differ.

This means that in general Z_L and Z'_L are not the same space, and to calculate Z_L we define a continuous family of finite reductions A_r^s, where s is a section in the bundle.

$$\Lambda\mathcal{L}(T(T^*L)) \to \Lambda T^*L.$$

Here $T(T^*L) \to T^*L$ is a symplectic bundle and $\mathcal{L}(T(T^*L)) \to T^*L$ is the associated fibration of Lagrangian Grassmanians. I.e. the fiber of the above fibration is $\Lambda\mathcal{L}(n) \simeq \Lambda(\mathrm{U}(n)/\mathrm{O}(n))$.

In the neighborhood of the zero section of L, we have two canonical sections of this bundle: ΛS_L, where S_L is the section in $\mathcal{L}(T(T^*L))$ which to a point associates the vertical directions w.r. to L, and similarly ΛS_N. The construction of A_r^s is such that the space Z_L does not change when perturbing s. This implies that if ΛS_L and ΛS_N where homotopic, we would in fact get that Z_L and Z'_L are homotopy equivalent. However, as mentioned before this is not the case in general[2].

If they are not homotopic we stabilize by adding trivial factors and get

$$A_r^{s\oplus\mathbb{R}^k} : T^*\Lambda_r(N \times \mathbb{R}^k) \cong T^*\Lambda_r N \times (\mathbb{R}^{2kn}, \omega_0) \to \mathbb{R}.$$

The space we get from our construction in this stabilized case is a suspension of Z_L. This is due to the fact that this function is just a constant quadratic term in the second variable $\mathbb{R}^{2kn}$. It is now a homotopy theoretical fact that: although ΛS_L and ΛS_N are not homotopic we can homotope $\Lambda S_N \oplus \mathbb{R}^k$ to take the form $\Lambda S_L \oplus J_*$, where $J_*\colon \Lambda T^*L \to \Lambda\mathcal{L}(k)$ is a curve of Lagrangians in $\mathbb{R}^{2n}$ dependant on the actual curve in the neighborhood of L. It is still true that $A_r^{\Lambda S_L \oplus J_*}$ is quadratic in the second variable, but this quadratic form is no longer the same for different choices of the first variable. We can now calculate that

$$\Sigma^l Z_L \simeq \mathrm{Th}(T\Lambda_r^{\mu_L}N \oplus \zeta)$$

where ζ is the negative eigen bundle of the quadratic form depending on J_*.

All in all we have that the collapse map is a map of Thom spaces

$$\mathrm{Th}(T\Lambda_r^{\mu}N \oplus \mathbb{R}^l) \simeq Z \to X/Y = Z_L \simeq \mathrm{Th}(T\Lambda_r^{\mu_L}L \oplus \zeta).$$

This was also noticed by Viterbo, except here we actually give an explicit discription of ζ. Using that L is Lagrangian, we get that $(\pi \circ j)^*TN \otimes \mathbb{C} \simeq TL \otimes \mathbb{C}$ and thus $2(\pi \circ j)^*TN \cong 2TL$. We can use this to remove all but one of the tangent spaces on both sides (a technical fact makes the above calculation true only for r odd). We get a map

---

[2] unless the nearby lagrangian conjecture is true, but we can of course not assume that.

$$\mathrm{Th}(\mathrm{Ev}_0^* TN \oplus \mathbb{R}^{l+l'}) \to \mathrm{Th}(\mathrm{Ev}_0^* TL \oplus \mathbb{R}^{l'} \oplus \zeta),$$

where Ev_0 is evaluation at the base point. This can also be written as

$$\Sigma^{l+l'}(\Lambda_r^{\mu} N)^{TN} \to \Sigma^{l'}(\Lambda_r^{\mu_L} L)^{TL\oplus\zeta}.$$

If we choose to to remove two extra copies of the tangent bundle on both sides and write $\zeta = \mathbb{R}^l \oplus \eta$, where η is a virtual bundle, we can write this as a map of Thom spectra:

$$(\Lambda_r^{\mu} N)^{-TN} \to (\Lambda_r^{\mu_L} L)^{-TL+\eta}.$$

Letting r and μ go to infinity we get the map

$$(\Lambda N)^{-TN} \to (\Lambda L)^{-TL+\eta}.$$

The reason why we write it like this is that the left hand side is a ring spectrum with the Chas-Sullivan product (see [3]), and in an opcomming paper we will prove that the right hand side is also a ring spectrum with at twisted version of the Chas-Sullivan product, and that this transfer is a ring spectrum homomorphism.

2 The Homotopy Index

This section is based on [4]. The theory has, however, been reformulated to suit the purpose of this paper.

Let M be a smooth open manifold, and let $f\colon M \to \mathbb{R}$ be a smooth function. A pseudo-gradient X for f is a smooth vector field on M such that the directional derivative $X(f)$ is positive at non-critical points and $X = 0$ at critical points. The choice of a pseudo-gradient is a contractible choice because a convex combination of pseudo-gradients is a pseudo-gradient. We will denote the flow of $-X$ by ψ_t.

Let a and b be regular values of f which are isolated from the critical values of f. We wish to define the *homotopy index* $I_a^b(f, X)$. In [4] this is called the Morse index, but we adopt the name homotopy index to avoid confusion.

Definition 2.1. An index pair (A, B) is a pair of subspaces of M satisfying the following properties

- I1: $B \subset A \subset f^{-1}([a, b])$.
- I2: A and B are compact.
- I3: $\mathrm{int}(A)$ contains all critical points of f with critical values in (a, b).
- I4: There is a function $c\colon A \to [0, \infty]$ such that for all $x \in A$ we have $\{t \geq 0 \mid \psi_t(x) \in A\} = [0, c(x)]$ and $\{t \mid \psi_t(x) \in B\} = \{c(x)\} \cap [0, \infty[$.

When such index pairs exist we define the Homotopy index

$$I_a^b(f, X) = [A/B],$$

where $[-]$ denotes homotopy equivalence class. If $X = \nabla f$ we write $I_a^b(f)$.

This is slightly different from [4], but we use this definition for simplicity. Note that it can be shown that the function c in I4 is continuous. B is called the *exit set*.

The following lemma shows that the homotopy index does not depend on the choice of index pair (A, B).

Lemma 2.2. *$I_a^b(f, X)$ is well-defined.*

Proof: If (A, B) and (A', B') is any pair of index pairs, we construct the intersection pair as follows

$$\begin{aligned} B'' &= (B \cup B') \cap (A \cap A') \\ A'' &= (A \cap A'). \end{aligned}$$

This is an index pair with $c''(x) = \min(c(x), c'(x))$, thus reducing to the case where A and B are subsets of A'.

Given any index pair (A, B), we can by using the flow ψ_t create new index pairs

$$\begin{aligned} A_2 &= \psi_t(A) \\ B_2 &= \psi_t(B) \end{aligned}$$

provided we stay within $f^{-1}([a, b])$. Being careful, one can choose t as a function of points in A and still get homeomorphic pairs. If B is not a subset of B', the function $c'_{|B}$ is not the zero function, and we can use the flow and the function c' to get the index pairs into a position where $B \subset B'$, thus reducing to $A \subset A'$ and $B \subset B'$.

Because $(A \cup B')/B' \simeq A/B$, we can replace (A, B) with $(A \cup B', B')$ which is a new index pair. This further reduces to the case $(A, B) \subset (A', B)$.

For any index pair (A, B) we can define a "flow" $P_t \colon A \to A$ by

$$P_t(x) = \varphi_{\min(t, c(x))}(x).$$

This map flows the set $\{x \mid c(x) \leq t\}$ into B. In particular it fixes B. This gives us a new pair $(P_t(A), B)$ with an equivalent quotient. In fact, the induced map

$$P_t \colon A/B \to A/B$$

is homotopic to the identity. In our case, the flow P_t for the small pair is the restriction of P'_t for the large pair.

Since $\overline{A' - A}$ is compact and $X_{|\overline{A'-A}}$ is non-zero, it has a lower bound, and we can thus find $t_0 > 0$ such that $P'_{t_0}(A') \subset A$. This defines a homotopy right inverse to the inclusion from A/B to A'/B. Doing the same for the pairs $(P'_{t_0}(A'), B) \subset (A, B)$ gives a homotopy right inverse to this map. □

We will need the concept of a ***good*** index pair (A, B). The definition uses a Riemannian metric on M, but does not depend on it.

Definition 2.3. An index pair (A, B) is called good if $B \subset f^{-1}(a)$ and if there is an $\varepsilon > 0$ such that when $\sup_{p \in A} \|X_p - X'_p\| < \varepsilon$ then the flow defined by $-X'$ exits A only through B.

This, together with an assumption on the critical points, will ensure that perturbing the data involved in defining the homotopy index does not change the homotopy index.

Lemma 2.4. *Let $f_s, s \in I$ be a homotopy of smooth functions, and $X_s, s \in I$ a homotopy of vector fields such that X_s is a pseudo-gradient for f_s. Let a, b be regular values isolated from the critical points for all f_s.*

Assume that for all $s_0 \in I$ we have a good index pair (A_{s_0}, B_{s_0}) defining $I_a^b(f_{s_0}, X_{s_0})$ and an $\varepsilon > 0$ such that $\operatorname{int}(A_{s_0})$ contains all critical points of f_s with critical value in (a,b) for $s \in [s_0 - \varepsilon, s_0 + \varepsilon]$. Then

$$I_a^b(f_0, X_0) = I_a^b(f_1, X_1).$$

Proof: Given s_0, we wish to prove that the good index pair $(A, B) = (A_{s_0}, B_{s_0})$ is an index pair for $I_a^b(f_s, X_s)$ when s is sufficiently close to s_0. However, this is not possible because we cannot be certain that I1 is satisfied. Because a and b are isolated critical points, we can replace a and b by $a-\delta$ and $b+\delta$ for some small δ without changing the indices. Now I1 is not a problem.

I2 is obvious and I3 is part of the assumptions in the lemma. Since the good pair assumption makes sure that we only exit A through B, we only need to prove that for any point x in B we have $\{t \geq 0 \mid \psi_t^s(x) \in A\} = \{0\}$ and I4 will follow. This is equivalent to proving that the flow does not return to A when exiting.

Since $-X_{s_0}(f_{s_0})$ restricted to B is negative, the same is true for $-X_s(f_{s_0})$ for s close to s_0, and thus we can find $\delta > 0$ such that $f_{s_0}(\psi_t^s(p))$ is strictly decreasing for $t \in [0, \delta]$ and $p \in B_{s_0}$. Because $f_{s_0}(B) = \{a\}$ and $f_{s_0}(A) \subset [a, b]$ we get for $p \in B$ and $t \in (0, \delta]$ that $\psi_t^s(p)$ is not in A and there is an $\varepsilon > 0$ such that $f_{s_0}(\psi_\delta^s(p)) < a - \varepsilon$. This implies (for s possibly closer to s_0) that $f_s(\psi_\delta^s(p)) < a - \varepsilon/2$. We can similarly assume that $f_s(A_{s_0}) > a - \varepsilon/3$, so the flow for $-X_s$ has exited B_{s_0} and will not return to A_{s_0} because it is a pseudo-gradient for f_s. □

As the following lemma shows, there is a way of producing good index pairs by using what we will call *cut-off* functions.

Lemma 2.5. *Assume that M has a Riemannian metric and that there exists non-negative functions $g_1, g_2, \ldots, g_n \colon M \to \mathbb{R}$ and constants $s_1 < t_1, s_2 < t_2, \ldots, s_n < t_n$ such that*

$$\begin{aligned} A &= f^{-1}([a,b]) \cap \{x \in M \mid g_j(x) \leq s_j + \frac{t_j - s_j}{b-a}(b - f(x))\} \\ B &= f^{-1}(a) \cap A \end{aligned}$$

are compact and the interior of A contains all the critical points of $f_{|(a,b)}$. If

$$-X_x(g_j) < \frac{t_j - s_j}{b-a} X_x(f) \tag{2}$$

for all $x \in \partial A$ which satisfy

$$g_j(x) = s_j + \frac{t_j - s_j}{b-a}(b - f(x)),$$

then (A, B) is a good index pair.

We will use this lemma repeatedly. In some cases when $X = \nabla f$ we will prove $\|\nabla g_j\| < \frac{t_j - s_j}{b-a}\|\nabla f\|$, which implies (2). Often (2) will be proven on much larger sets than needed.

Proof: At any point $x \in \partial A$ we must have $f(x) - b \leq 0$ and

$$g_j(x) - s_j - \frac{t_j - s_j}{b-a}(b - f(x)) \leq 0$$

satisfied. The assumptions in the lemma and the fact that X is a pseudo-gradient ensure that for any vector v close to X_x we have: if any of these inequalities is an equality then the directional derivatives of the left hand side in direction $-v$ is negative. So $-v$ points into A, except if the equality $f(x) = a$ is satisfied in which case $-v$ must point out of the set. The boundary is compact so there is an $\varepsilon > 0$ such that this holds for all v and all x if $\|v - X_x\| < \varepsilon$ $\square$

Some very important aspects of homotopy indices are the natural inclusion and quotient maps

$$i\colon I_a^b(f,X) \to I_a^c(f,X)$$
$$q\colon I_a^c(f,X) \to I_b^c(f,x)$$

where $a < b < c$. These maps are constructed as follows. If (A,B) is an index pair for $I_a^c(f,X)$ then $(A \cap f^{-1}([a,b]), B \cap f^{-1}([a,b]))$ is an index pair for $I_a^b(f,X)$, and the inclusion is the obvious one. Similarly, $(A \cap f^{-1}([b,c]), A \cap f^{-1}(\{b\}) \cup B \cap f^{-1}([b,c]))$ is an index pair for $I_b^c(f,X)$, and the quotient is the map collapsing the set $(A \cap f^{-1}([a,b]))/B$.

3 The Action Integral in Cotangent Bundles

All parts of this section are well-known, but the methods are vital to the construction, and we need to introduce the notation anyway.

Let N be a smooth n-dimensional manifold. We denote points in the cotangent bundle T^*N by (q,p), where q is in N and p is a cotangent vector. Let $\pi\colon T^*N \to N$ be the projection onto the base and define the canonical 1-form $\lambda \in \Omega^1(T^*N)$ and 2-form $\omega \in \Omega^2(T^*N)$ by

$$\lambda_{q,p}(v) = p(\pi_*(v))\omega \qquad = d\lambda.$$

The form ω is non-degenerate and thus defines a canonical symplectic structure on T^*M.

Given any smooth Hamiltonian $H\colon T^*N \to \mathbb{R}$, we can define the associated Hamiltonian vector field X_H by the formula $dH = \omega(X_H, -)$. This is well-defined because ω is non-degenerate. The flow of X_H will be denoted φ_t and is called the Hamiltonian flow.

Given any metric on N we can induce a metric on T^*N in the following way: at each point (q,p) we split the tangent space $T_{(q,p)}(T^*N)$ in two components, the vertical, which is canonically defined without the metric as the fiber directions, and the horizontal defined by the connection given by the metric on N. This identifies $T_{(q,p)}T^*N$ with $T_qN \oplus T_q^*N$, on which we use the metric from M to define the inner product on each factor - making this splitting orthogonal. We can also define an almost complex structure J in this splitting by using the isometry $\phi_q \colon T_qN \to T_q^*N$ induced by the metric on N

$$J(\delta q, \delta p) = (-\phi^{-1}(\delta p), \phi(\delta q)).$$

This is compatible with the symplectic structure and the induced metric. The formula for X_H can be rewritten using this metric and almost complex structure as

$$X_H = -J\nabla H. \tag{3}$$

Let ΛM be the space of piecewise smooth and continuous maps from S^1 to M. The action $A_H \colon \Lambda T^*N \to \mathbb{R}$ is defined by

$$A_H(\gamma) = \int_\gamma \lambda - \int_{S^1} H(\gamma(t))dt = \int_\gamma (\lambda - Hdt).$$

It is known that the critical points of this integral are precisely the 1-periodic orbits of the Hamiltonian flow (a calculation in the following section proves this).

For a moment we look at the special case in which H only depends on the length of the cotangent vector - that is

$$H(q,p) = h(\|p\|),$$

where $h \colon \mathbb{R} \to \mathbb{R}$. For H to be smooth there are some restriction on the deriviatives of h at 0.

In this case we calculate the gradient of H in the orthogonal splitting:

$$\nabla H = (0, h'(\|p\|)p)$$

We get 0 in the first factor because parallel transport does not change the norm of p. Using equation (3) we see that

$$X_H = -J(0, h'(\|p\|)p) = (h'(\|p\|)\phi^{-1}(p), 0) = (h'(\|p\|)p, 0).$$

As hinted, from now on we will suppress ϕ from the notation.

Remark 3.1. Because this vector field is 0 on the last factor, it will parallel transport p and hence this becomes a reparametrization of the geodesic flow on N. This describes the 1-periodic orbits as closed geodesics on N with lengths corresponding to $h'(\|p\|)$. The action of these orbits is calculated to be $\|p\|h'(\|p\|) - h(\|p\|)$. This corresponds to taking minus the intersection of the y-axis with the tangent of h at the point $(x, h(x))$ as in figure 1. This geometric formula for calculating the action is very useful for this type of Hamiltonian, and will be used repeatedly.

4 Finite Reductions of the Action Integral in Cotangent Bundles for a Hamiltonian Linear at Infinity

This section is inspired by work in [10] and [2], which use the broken geodesic approach to do what Floer homology later did more generally. We will define



Figure 1: Geometric calculation of critical values.

finite dimensional reductions of the action integral in cotangent bundles, and compute their homotopy indices. This is the approach used by Viterbo to construct his transfer in [10]. He uses the theory of generating functions, but we will more explicitly construct the functions and obtain more control over the homotopy indices.

We assume that N is a closed manifold with a Riemannian structure and an injective radius $2\varepsilon_0$. On T^*N we have the induced metric. We also assume that $H\colon T^*N \to \mathbb{R}$ is a smooth Hamiltonian with the following property: there exist $R, \mu, c \in \mathbb{R}$, μ not a geodesic length, such that $H(q,p) = \mu\|p\| + c$ when $\|p\| > R$. The Hamiltonian flow of H is denoted φ_t.

We assume that $C_1 \geq \mu$ is an upper bound for $\|\nabla H\|$ on all of T^*N, and C_2 is an upper bound for the norm of the covariant Hessian $\|\nabla^2 H\|$ on the set $T^*_R N = \{(q,p) \in T^*N \mid \|p\| \leq R\}$. We define the space of piecewise geodesics as

$$\Lambda_r N = \{(q_j)_{j\in\mathbb{Z}_r} \in N^r \mid \operatorname{dist}(q_j, q_{j+1}) < \varepsilon_0\},$$

where $\operatorname{dist}(-,-)$ is the distance in N. This implies

$$T^*\Lambda_r N = \{(q_j, p_j)_{j\in\mathbb{Z}_r} \in (T^*N)^r \mid \operatorname{dist}(q_j, q_{j+1}) < \varepsilon_0\}.$$

We will denote a point in this space by $\overrightarrow{z} = (\overrightarrow{q}, \overrightarrow{p})$, and a single coordinate by $z_j = (q_j, p_j) \in T^*N$. These two spaces are given the product Riemannian structures from N and T^*N.

We will define functions resembling A_H on $T^*\Lambda_r N$ having the same critical points (the 1-periodic orbits) with the same critical values, and we will prove that for r large enough these functions admit good index pairs. In section 5 we will define generalizations of these, and in remark 5.5 we explain how they can be constructed using embeddings $i_r\colon T^*\Lambda_r N \to \Lambda N$ as explained in the overview in the introduction.

Definition 4.1. For $C_1/r < \varepsilon_0/3$ we define

$$A_r(\overrightarrow{z}) = \sum_{j\in\mathbb{Z}_r} \Big(\int_{\gamma_j} \lambda - H dt\Big) + \sum_{j\in\mathbb{Z}_r} p_j^- \varepsilon_{q_j},$$

where $\gamma_j\colon [0,1/r] \to T^*N$ is the curve $\gamma_j(t) = \varphi_t(q_j, p_j)$,

$$(q_j^-, p_j^-) = \varphi_{1/r}(q_{j-1}, p_{j-1}) \qquad \text{and} \qquad \varepsilon_{q_j} = \exp^{-1}_{q_j^-}(q_j) \in T_{q_j^-}N,$$

with $\exp\colon TN \to N$ the exponential map.

The term $p_j^- \varepsilon_{q_j}$ is the pairing of cotangent vectors with tangent vectors. This function is well-defined because $C_1/r < \varepsilon_0$ implies that the distance between q_j^- and q_j is less than $2\varepsilon_0$. It is for later reasons that we assume the stricter inequality $C_1/r < \varepsilon_0/3$. Notice that if $\|p_j\| > R$, then p_{j+1}^- is the parallel transport of p_j by a geodesic in the direction of p_j, and thus $\|p_{j+1}^-\| = \|p_j\|$ (see the previous section for details).

When evaluating the function A_r on an r-pieced dissection of a 1-periodic orbit, we see that the last term vanishes because the γ_j's fit together to a closed curve. So A_r equals the action integral on such a curve.

Before the next lemma we need a few more definitions and a few abbreviations. Consider the commutative diagram of isomorphisms

$$\begin{tikzcd}
T_qN \arrow[r, "P_{q,q'}"] \arrow[d, "\phi"] & T_{q'}N \arrow[d, "\phi"] \\
T_q^*N \arrow[r, "P^*_{q,q'}"] & T^*_{q'}N.
\end{tikzcd}$$

The isomorphism ϕ is the one induced by the metric, which we suppressed from the notation in the previous section. We will do so again. The functions $P_{q,q'} = P^*_{q,q'}$ are parallel transport along the unique geodesic connecting q and q' when $\operatorname{dist}(q,q') < 2\varepsilon_0$. We use P_{q_j,q_j^-} to define

$$\varepsilon_{p_j} = P_{q_j,q_j^-}(p_j) - p_j^- \in T^*_{q_j^-}N \simeq T_{q_j^-}N$$

both ε_{q_j} and ε_{p_j} are vectors in the (co)tangent space at q_j^-. We can parallel transport any vector in the (co)tangent space at q_j^- to the (co)tangent space at q_j and q_{j-1}. The resulting vectors will by further abuse of notation be denoted the same. So e.g. for ε_{p_j} this means

$$\varepsilon_{p_j} = p_j - p_j^- = p_j - P_{q_j^-,q_j}(p_j^-) \in T_{q_j}N$$

and

$$\varepsilon_{p_j} = p_j - p_j^- = P_{q_j^-,q_{j-1}}\big(p_j - P_{q_j,q_j^-}(p_j^-)\big) \in T^*_{q_{j+1}}N.$$

We also define $P = \max_j(\|p_j\|)$.

Lemma 4.2. *There exists a constant $K' > 0$ which is independent of r, C_1 and C_2 such that for $K = K'(C_2 + C_1)$ we have*

$$\begin{aligned}
\|\nabla_{q_j} A_r + \varepsilon_{p_j}\| &\leq K \max(R,P)(\|\varepsilon_{q_j}\| + \|\varepsilon_{q_{j+1}}\|) \\
\|\nabla_{p_j} A_r - \varepsilon_{q_{j+1}}\| &\leq \frac{K}{r}\|\varepsilon_{q_{j+1}}\|,
\end{aligned}$$

where $\nabla_{q_j} A_r \oplus \nabla_{p_j} A_r = \nabla_{z_j} A_r$ is the gradient with respect to the jth component in $T^*\Lambda_r N$. Furthermore, for $r > K$ the only critical points of A_r are the r-pieced dissections of the 1-periodic flow curves for the Hamiltonian flow.

Note that the r-pieced dissections of the 1-periodic orbits correspond to the points where $\varepsilon_{q_j} = \varepsilon_{p_j} = 0$ for all j.

Proof: We start out by considering one of the integration terms. In the following $\delta\gamma_j$ is a variation of the curve γ_j (a tangent field along γ_j).

$$\begin{aligned}\nabla(\int_{\gamma_j} & \lambda - H dt)(\delta\gamma_j) \\ &= \int_0^{1/r} \gamma_{jp}(\nabla_t \delta\gamma_{jq}) + (\delta\gamma_{jp})\gamma'_{jq} - \nabla_{\gamma_j(t)} H(\delta\gamma_j(t))dt \\ &= [\gamma_{jp}(t)\delta\gamma_{jq}(t)]_0^{1/r} + \int_0^{1/r} -\gamma'_{jp}(\delta\gamma_{jq}) + (\delta\gamma_{jp})\gamma'_{jq} - \nabla_{\gamma_j(t)} H(\delta\gamma_j(t))dt \\ &= -p_j\delta q_j + p^-_{j+1}\delta q^-_{j+1} - \int_0^{1/r} (J\gamma'_j(t) + \nabla_{\gamma_j(t)} H)(\delta\gamma_j(t))dt \qquad (4) \\ &= -p_j\delta q_j + p^-_{j+1}\delta q^-_{j+1}\end{aligned}$$

The integral vanishes because γ_j is a flow curve, and thus $\gamma'_j(t) = -J\nabla_{\gamma_j(t)} H$. This is a standard calculation, and it is also a proof that the 1-periodic orbits of the flow X_H are the critical points of the action integral.

Our function depends on $(q_j, p_j), j \in \mathbb{Z}_r$, but it is convenient to continue for a while calculating the gradient as if the function depended on both $(q_j, p_j)_{j\in\mathbb{Z}_r}$ and $(q^-_j, p^-_j)_{j\in\mathbb{Z}_r}$ as independent variables. Formally we define

$$T^*\Lambda'_r N = \{(q_j, p_j, q^-_j, p^-_j) \in T^*N^{2r} \mid \operatorname{dist}(q_j, q^-_j) < 2\varepsilon_0, \operatorname{dist}(q^-_j, q_{j+1}) < 2\varepsilon_0\}.$$

We have an embedding

$$\iota\colon T^*\Lambda_r N \to T^*\Lambda'_r N$$

defined by setting $(q^-_j, p^-_j) = \varphi_{1/r}(q_{j-1}, p_{j-1})$ for all j.

We wish to extend the function A_r to a function A'_r defined on $T^*\Lambda'_r N$, and calculate the gradient of A'_r on the image of ι. So, for each point (q, p, q^-, p^-) in T^*N^2 with $\operatorname{dist}(q, q^-) \le 2\varepsilon_0$, we choose a curve connecting (q, p) with (q^-, p^-), such that if $\varphi_{1/r}(q, p) = (q^-, p^-)$ then we chose the flow curve $\varphi_t(q, p)$ used in the definition of A_r. By integrating over the chosen curve, we extend the definition of the integration term of A_r to $T^*\Lambda'_r N$. The calculation above shows that the gradient of this term on the image of ι does *not* depend on the choice of curves, and we have in fact already calculated it.

The second term we extend simply by using the same expression

$$\sum_j p^-_j \exp^{-1}_{q^-_j}(q_j).$$

So we need to calculate the gradient of the function

$$f(q, q^-, p^-) = p^- \exp^{-1}_{q^-}(q) = p^-\varepsilon_q, \qquad q \in N, (q^-, p^-) \in T^*N.$$

One of the components is obvious:

$$\nabla_{p^-} f = \varepsilon_q,$$

however, when moving q and q^- we get some interference from the curvature of N. Assume that $\|p^-\| = 1$. If $\varepsilon_q = 0$ we see that

$$\begin{aligned}\nabla_{q^-} f &= -p^- \\ \nabla_q f &= p^-.\end{aligned}$$

So by compactness we can find $k > 0$ such that

$$\begin{aligned} \|\nabla_{q^-} f + p^-\| &\leq k\|\varepsilon_q\| \\ \|\nabla_q f - p^-\| &\leq k\|\varepsilon_q\|. \end{aligned}$$

Here we have used the abuse of notation discussed just before the lemma to define p^- as a tangent vector at q. Using that $f(q, q^-, sp^-) = sf(q, q^-, p^-)$ we get a bound in the general case:

$$\begin{aligned} \|\nabla_{q^-} f + p^-\| \leq& k\|p^-\|\|\varepsilon_q\| \\ \|\nabla_q f - p^-\| \leq& k\|p^-\|\|\varepsilon_q\|. \end{aligned}$$

Adding the two terms of the gradients we obtain

$$\nabla_{q_j, p_j, q_j^-, p_j^-} A'_r = (-\varepsilon_{p_j} + \mathrm{bo}(k\|p_j^-\|\|\varepsilon_{q_j}\|), 0, \mathrm{bo}(k\|p_j^-\|\|\varepsilon_{q_j}\|), \varepsilon_{q_j}), \tag{5}$$

where the notation $\mathrm{bo}(c)$ means some term bounded by c, i.e. $a = b + \mathrm{bo}(c)$ is equivalent to $\|a - b\| \leq c$.

If $H = 0$ we have $(q_j^-, p_j^-) = (q_{j-1}, p_{j-1})$, and the gradient is just the sum of the two components:

$$\nabla_{q_j, p_j} A_r = \big(-\varepsilon_{p_j} + \mathrm{bo}(k(\|p_j^-\|\|\varepsilon_{q_j}\| + \|p_{j+1}^-\|\|\varepsilon_{q_{j+1}}\|)), \varepsilon_{q_{j+1}}\big).$$

However, if H is non-zero we need to understand the differential of the function $\varphi_{1/r}$, and use that for $v \in T_{q_j, p_j}(T^*N)$

$$v(A_r) = (D_{q_j, p_j}(\mathrm{Id} \times \varphi_{1/r})(v))(A'_r). \tag{6}$$

Assume $(q^-, p^-) = \varphi_{1/r}(q, p)$. We would like to compare $D_{q,p}\varphi_{1/r}$ to the parallel transport of (co)tangent vectors from q to q^- used in defining $\varepsilon_{q_{j+1}}$ at $T_{q_j}N$. That is, by using the splittings

$$T_{q,p}T^*N = T_qN \oplus T_q^*N, \quad T_{q^-, p^-}T^*N = T_{q^-}N \oplus T_{q^-}^*N$$

and the parallel transports we define

$$T_{q,p} = P_{q,q^-} \oplus P_{q,q^-}^* : T_{q,p}T^*N \to T_{q^-, p^-}T^*N.$$

First consider the compact set T_R^*N. On this set we have $\|D_{q,p}\varphi_{1/r} - T_{q,p}\| < k_2/r$ for some k_2. The question is how k_2 depends on C_1 and C_2. Take a normal neighborhood of (q, p) (this identifies the tangent spaces locally). In this, $T_{q,p}$ is C_1k_3/r close to the identity because the length of the flow curve is less than C_1/r, and therefore k_3 only depends on the metric on the compact set $\|p\| \leq R$. On the other hand, we have $D_{q,p}\varphi_{1/r}$ is also close to the identity, but how close depends on the first order derivatives of $X_H = -J\nabla H$, which is the vector field defining the flow φ_t. This is bounded in the normal chart by $c_1C_1 + c_2C_2$, for some c_2 depending on the metric and some c_1 depending on how parallel the complex structure is. We conclude that k_2 can be chosen as some constant times $(C_1 + C_2)$.

On the rest of the cotangent bundle we will get a bound using the action of $\mathbb{R}_+$ given by $t(q, p) = (q, tp)$ and the fact that

$$T_{q,p} = T_{q,tp} \qquad \text{and} \qquad \varphi_{1/r}(q, tp) = t\varphi_{1/r}(q, p).$$

The latter only applies when $\|p\| \geq R$ and $\|tp\| \geq R$, because this is where H has the special form $\mu\|p\| + c$. Let $a_t(q,p) = (q, tp)$. Then the differential of a_t in (q,p) splitting is

$$D_{q,p}a_t = \begin{bmatrix} \mathrm{Id} & 0 \\ 0 & t\,\mathrm{Id} \end{bmatrix}.$$

So by decomposing

$$\varphi_{1/r}(q,p) = (\|p\|/R)\varphi_{1/r}(q,(R/\|p\|)p) = a_{\|p\|/R} \circ \varphi_{1/r} \circ a_{R/\|p\|}$$

whenever $\|p\| > R$, and using the bound we already have on the compact set $p \leq R$, we get the bound

$$D_{q,p}\varphi_{1/r} - T_{q,p} = \begin{bmatrix} \mathrm{bo}(\frac{k_2}{r}) & \mathrm{bo}(\frac{k_2 R}{r\max(R,\|p\|)}) \\ \mathrm{bo}(\frac{k_2\max(R,\|p\|)}{rR}) & \mathrm{bo}(\frac{k_2}{r}) \end{bmatrix} \tag{7}$$

for all $(q,p) \in T^*N$. Using equations (5), (6) and (7) together with

$$\frac{\|p^-_{j+1}\|}{\max(R,\|p_j\|)} \leq 1,$$

we get the estimates

$$\begin{aligned} \nabla_{q_j}A_r &= \Big(-\varepsilon_{p_j} + \mathrm{bo}\big(K\max(R,P)(\|\varepsilon_{q_j}\| + \|\varepsilon_{q_{j+1}}\|)\big)\Big) \\ \nabla_{p_j}A_r &= \Big(\varepsilon_{q_{j+1}} + \mathrm{bo}(\frac{K}{r}\|\varepsilon_{q_{j+1}}\|)\Big) \end{aligned}$$

with K some real number proportional to k_2. This proves the first part of the lemma.

The last part of the lemma is simple: assume $\frac{K}{r} < 1$, then $\nabla_{p_j}A_r$ is zero if and only if $\varepsilon_{q_{j+1}}$ is zero, and having that for all j we can conclude the same for $\nabla_{q_j}A_r$ and ε_{p_j}. So the critical points are exactly the ones where $\|\varepsilon_{q_j}\| = \|\varepsilon_{p_j}\| = 0$ for all j. □

The function A_r with its gradient does not necessarily have index pairs, but we define a pseudo-gradient X with which it does. On the set where $\max_j\|\varepsilon_{q_j}\| < \varepsilon_0/4$ we use the gradient of A_r, and on the set $\max_j\|\varepsilon_{q_j}\| > \varepsilon_0/3$ we keep the non-zero p-component of the gradient of A_r, but use 0 as the q-component. In between we use some convex combination of them. So

$$X \cdot \nabla A_r \geq \|X\|^2 \geq \sum_j \|\nabla_{p_j}A_r\|^2, \tag{8}$$

and as we only made X different from the gradient on a set where this is non-zero, it is indeed as pseudo-gradient.

Lemma 4.3. *Let a and b be regular values of A_r, and let K be as in the previous lemma. If $r > K$ then (A_r, X) has a good index pair, which will contain all the critical points of any A_r coming from a small C^1-perturbation of H fixing H on the set where $\|p\| > R$. Furthermore, if X' is another pseudo-gradient such that $X \neq X'$ is contained in something compact, it will not change the homotopy index.*

Proof: We will need a global lower bound for $\|X\|$ on the set where $P > 2R$. First we prove this on the set where $X \neq \nabla A_r$. Here we have some $\|\varepsilon_{q_j}\| > \varepsilon_0/4$, and from the previous lemma we have $\|X\| > (1-K/r)\|\varepsilon_{q_j}\| > (1-K/r)\varepsilon_0/4$.

So now we look at the case where $X = \nabla A_r$. We will get a lower bound if we show that there exist $k_1, k_2 \in \mathbb{R}_+$ such that if $G_p = \sum_j \|\nabla_{p_j} A_r\| < k_1$, then $G_q = \sum_j \|\nabla_{q_j} A_r\| > k_2$. Define

$$L_q = \sum_j \|\varepsilon_{q_j}\| \qquad \text{and} \qquad L_p = \sum_j \|\varepsilon_{p_j}\|.$$

The statement can, because of the approximation of $\nabla_{p_j} A_r$ in the previous lemma, be reduced to: there exist k_1, k_2 such that $L_q < k_1$ implies $G_q > k_2$. This is the statement we will prove.

Define $\underline{P} = \min_j \|p_j\|$. There are no 1-periodic flow curves on the compact set $R \leq \|p\| \leq 2R$, so there must exist $0 < c < 1$ such that $L_q + L_p > c$ for curves with all z_j's contained in this set. Claim: $k_1 = \min(c/2, \frac{1}{4K}, \frac{c}{8KR})$ works. We will divide the proof of this claim into two cases.

First case: $\underline{P} < P/2$. We know that for some j we have $\|p_j\| = P \geq 2R$ and for some j' we have $\|p_{j'}\| < P/2$. The "curve" $\overrightarrow{z}$ has to move this distance in p-direction and back again. So because $|\|p_j\| - \|p_{j-1}\|| = |\|p_j\| - \|p_j^-\|| < \|\varepsilon_{p_j}\|$ when $\|p_j\| \geq R$, we get that $L_p > P$. With the bound $L_q < 1/(4K)$ we see that this implies

$$\begin{aligned} G_q &> \sum_j \|-\varepsilon_{p_j} + \mathrm{bo}\big(KP(\|\varepsilon_{q_j}\| + \|\varepsilon_{q_{j+1}}\|)\big)\| \\ &> \sum_j \|\varepsilon_{p_j}\| - KP(\|\varepsilon_{q_j}\| + \|\varepsilon_{q_{j+1}}\|) > (P - \frac{P}{2}) > R, \end{aligned}$$

which is a positive constant.

The second case: $\underline{P} \geq P/2$. In this case we can, because the flow is equivariant with respect to the $\mathbb{R}_+$ action on the set $\|p\| \geq R$, multiply our "piecewise flow curve" with $2R/P$ to obtain a piecewise flow curve on the compact set $R \leq \|p\| \leq 2R$. This does not change any of the ε_{q_j}'s, but it scales the ε_{p_j}'s so we can conclude that the original curve satisfies

$$\frac{2R}{P} L_p + L_q > c.$$

Because $L_q < c/2$ this implies that $L_p > \frac{cP}{4R}$, which implies by using the bound $L_q < \frac{c}{8KR}$ that

$$G_q > \sum_j \|\varepsilon_{p_j}\| - KP(\|\varepsilon_{q_j}\| + \|\varepsilon_{q_{j+1}}\|) > \frac{cP}{4R} - \frac{cP}{8R} > \frac{c}{4}.$$

This is again a positive constant.

So we have proved that there exists $C > 0$ such that

$$\|X\| > C$$

on the non-compact set where $P > 2R$.

Because of equation (8) we obtain

$$\big|X \cdot \nabla \|p_j\|\big| \leq \|X\| \leq C^{-1}\|X\|^2 \leq C^{-1}X \cdot \nabla A_r.$$

So we can use $\|p_j\|$ as a cut-off function with $s_j > 2R$ and $t_j - s_j > C^{-1}(b-a)$ (see lemma 2.5).

Define $f_j(\overrightarrow{z}) = \mathrm{dist}(q_j, q_{j+1})$. If $f_j < 2\varepsilon_0/3$ we will, because of the assumption $C_1/r < \varepsilon_0/3$ in definition 4.1, get $\|\varepsilon_{q_j}\| > \varepsilon_0/3$. The way we defined X is such that at a point like this we have $X \cdot \nabla f_j = 0$. So we can use f_j as a cut-off function with any $\varepsilon_0 > t_j > s_j > 2\varepsilon_0/3$. We now have enough cut-off functions to ensure that when using lemma 2.5 we get a compact index pair inside our open manifold.

The fact that our pair contains the critical points of A_r when H is perturbed follows from: as long as C_1/r is less that $\varepsilon_0/3$ we can only get critical points when $f_j < \varepsilon_0/3$, and we get no critical points with $p_j > R$ because of the way we defined H on $\|p\| > R$.

The last statement follows because any compact subset can be contained in an index pair constructed in this way, and changing the pseudo-gradient on the interior of an index pair will not change the homotopy index. ☐

Because all critical points of A_r lie in the compact set $P < R$, the set of critical values must be compact. So the total index $I(A_r, X)$ is well-defined. The lemma above tells us that changing the Hamiltonian within the proper bounds and conditions does not change the total index. So we define a specific Hamiltonian and calculate the total index. To do this we first need part of the following small lemma.

Lemma 4.4. *The set of lengths of geodesics in N is closed and with measure zero.*

Proof: For any c there is a compact manifold inside the space of curves containing all geodesics with energy less than c. On this manifold the energy is smooth, and has the geodesics as critical points, so by Sards theorem the energy spectrum below c of geodesics is closed and with measure zero. But since c is arbitrary this is true for the entire spectrum. For geodesics the length is easily related to the energy. So we conclude that the length spectrum is closed and with measure zero. ☐

Define $H_0(q,p) = h(\|p\|)$ where $h(t) = \frac{\mu}{2}t^2$ when $t < \frac{\mu-\varepsilon}{\mu}$ for some $\varepsilon > 0$ such that $[\mu-\varepsilon, \mu]$ does not contain any geodesic lengths.

We still need $h(t) = \mu t + c$ outside a compact set, but we also want h to be convex so that all the 1-periodic orbits will lie in the set where h is quadratic. In fact we want h'' to be a bump function constantly equal to μ on the set $\|p\| < \frac{\mu-\varepsilon}{\mu}$ and zero on the set $\|p\| > \frac{\mu+\varepsilon}{\mu}$ such that it integrates to μ. Notice that $\frac{\mu+\varepsilon}{\mu} < 2$, so for this Hamiltonian we can choose $R = 2$.

Proposition 4.5. *There exists a constant $C > 0$ such that for any Hamiltonian H_0 described above we have that $r > C\mu$ implies existence of index pairs and*

$$I(A_r, X) = \mathrm{Th}(T\Lambda_r^{\mu}N),$$

where $\Lambda^\mu_r N$ *is the manifold of piecewise geodesic curves in* N*, each piece having length less than* μ/r*.*

Proof: We define $\overline{\Lambda}^a_r N$ to be piecewise geodesics, each piece having length less than or equal to a/r. We then define an embedding

$$i\colon \overline{\Lambda}^{(\mu-\varepsilon)}_r N \to T^*\Lambda_r N,$$

where the ε and μ comes from the definition of H_0 above and i is given by

$$(i(\overrightarrow{q}))_j = \big(q_j, \mu^{-1} r \exp^{-1}_{q_j}(q_{j+1})\big).$$

This is a section in the bundle $T^*\Lambda_r N \to \Lambda_r N$ restricted to $\overline{\Lambda}^{(\mu-\varepsilon)}_r N$. Because $\|\mu^{-1} r \exp^{-1}_{q_j}(q_{j+1})\| < \frac{\mu-\varepsilon}{\mu}$, the point $(i(\overrightarrow{q}))_j$ will lie in the set where h is quadratic. This means that the flow curve γ_j is easy to understand, and in fact we have chosen p_j in such a way that $q^-_{j+1} = q_{j+1}$. This implies that on the image of i all ε_{q_j} are 0. In fact, this is the unique point in the fiber over q_j that flows to the fiber over q_{j+1}. The image of this embedding contains all the critical points of A_r, because it contains all the curves with $\|p_j\| \le \mu - \varepsilon$ and $\varepsilon_{q_j} = 0$.

We will use the fiber directions ($\overrightarrow{p}$ directions) as a normal bundle. In fact because $\varepsilon_{q_j} = 0$ for all j, lemma 4.2 tells us that $\nabla_{p_j} A_r = 0$, and because this is the only point in the fiber that flows to a point over q_{j+1}, this is the only critical point when restricting A_r to the fiber.

Claim: if $\overrightarrow{q} \in \overline{\Lambda}^{\mu-\varepsilon}_r N$ is fixed then the function $A_r(\overrightarrow{q}, \overrightarrow{p})$ goes to $-\infty$ as $\|\overrightarrow{p}\|$ goes to ∞.

This makes the point of the embedding the global maximum in the fiber.

Proof of claim: the condition $\|\overrightarrow{p}\| \to \infty$ is equivalent to $\|p_j\| \to \infty$ for some j. So we look at the terms in the definition of our finite reduction that involves p_j:

$$f(p_j) = \int_{\gamma_j} (\lambda - H dt) + p^-_{j+1} \varepsilon_{q_{j+}}.$$

Assume that $\|p_j\| > 2$. The integration part was already calculated in the previous section and is $(\|p_j\| h'(\|p_j\|) - h(\|p_j\|))/r$, which is constant on the set $\|p_j\| > 2$. Because $\operatorname{dist}(q_j, q_{j+1}) \le (\mu - \varepsilon)/r$ and $\operatorname{dist}(q_j, q^-_{j+1}) = \mu/r$, we are in the situation depicted in figure 2. Take the metric we have on N and multiply it with r/μ, and take a normal chart around q_j in this new metric. Then the circle in the picture is mapped to the unit circle. If the metric on the unit disc is "flat enough" then the paring $p^-_j \varepsilon_{q_{j+1}}$ will be negative, so by making μ/r smaller than some constant depending on the metric, we know that the second term is negative when $p_j > 2$. However, multiplying p_j with a constant larger than 1 scales this term by the same constant. So, A_r goes to $-\infty$ if $\|p_j\|$ goes to ∞.

On the image of the embedding the last term vanishes, and

$$A_r(i(\overrightarrow{q})) = \sum_j \int_{\gamma_j} (p dq - H dt) = \frac{1}{2\mu} \sum_j r \|\exp^{-1}_{q_j}(q_{j+1})\|^2.$$



Figure 2: Position of points when the norm of p_j is larger than 2.

This is μ^{-1} times the energy functional

$$e(\gamma) = \frac{1}{2}\int_0^1 \|\gamma'(t)\| dt$$

evaluated on the piecewise geodesic $\overrightarrow{q}$. This is positive and we conclude that if we look at the set defined by $A_r \geq -1$ intersected with one of the fibers, we get a bounded set diffeomorphic to a closed disc. This is true over every point in the compact set $\overline{\Lambda}_r^{(\mu-\varepsilon)}N$, so the set

$$A = \{(\overrightarrow{q}, \overrightarrow{p}) \mid \overrightarrow{q} \in \overline{\Lambda}_r^{(\mu-\varepsilon)}N, A_r(\overrightarrow{q}, \overrightarrow{p}) \geq -1\}$$

is compact and has points in each fiber.

We now change the pseudo-gradient X to another X'. We do this on a neighborhood of A - say the interior of the compact set

$$A' = \{(\overrightarrow{q}, \overrightarrow{p}) \mid \overrightarrow{q} \in \Lambda_r^{(\mu-\varepsilon/2)}N, A_r(\overrightarrow{q}, \overrightarrow{p}) < -2\}.$$

Lemma 4.3 tells us that this does not change the homotopy index. We will only specify X' on A, because the choice of a pseudo-gradient is contractible, and so it is easy to extend and interpolate. As we noted before, the $\overrightarrow{p}$ part of the gradient of A_r is non-zero except on the embedding, so we will use this as X' except in a small neighborhood of the embedding. On this set we will use the gradient of the energy functional, which we proved coincided with A_r on the embedding. Since minus the gradient of the energy functional on piecewise geodesics defines a flow that flows in a direction in which the longest geodesic piece gets shorter or stays the same length, we know that the gradient of the energy will preserve $\overline{\Lambda}_r^{(\varepsilon-\mu)}N$. This implies that the pseudo-gradient X' will point into or be parallel to the boundary of A close to the embedding. So the points at which the flow of $-X'$ exits A is precisely the points where $A_r = -1$. In each fiber this is the boundary of the disc defined by $A_r \geq -1$. So if we define

$$B = \{(\overrightarrow{q}, \overrightarrow{p}) \mid \overrightarrow{q} \in \overline{\Lambda}_r^{(\mu-\varepsilon)}N, A_r(\overrightarrow{q}, \overrightarrow{p}) = -1\},$$

then (A, B) is an index pair for $I(A_r, X')$, and they are the appropriate disc and sphere bundles needed to prove the lemma. □

This proof can be slightly modified to prove that the homotopy index of this specific Hamiltonian with respect to an interval (a, b) (simply by using the same argument for the embedding restricted to an appropriate subspace) is

$$I_a^b(A_r, X) = \mathrm{Th}(T\Lambda_r^{\min(\sqrt{2\mu a},\mu)}N)\Big/\mathrm{Th}(\Lambda_r^{\min(\sqrt{2\mu b},\mu)}N).$$

Here the $x \mapsto \sqrt{2\mu x}$ is the conversion from μ^{-1} times energy to length. This is needed because the critical value corresponding to a geodesic was calculated in the proof to be μ^{-1} times the energy.

It seems that increasing r by 1 gives a Thom suspension of the total homotopy index. The next lemma proves this for any a, b and H.

Lemma 4.6. *Assume that $r > K$ (from lemma 4.3). The index $I_a^b(A_{r+1}, X_{r+1})$ is the relative Thom suspension of $I_a^b(A_r, X_r)$ by the bundle*

$$\mathrm{Ev}_{q_0}^* TN,$$

where $\mathrm{Ev}_{q_0} : T^*\Lambda_r N \to N$ is the function defined by $\mathrm{Ev}_{q_0}(\overrightarrow{z}) = q_0$.

Note that the relative Thom suspension of A/B with a bundle ζ over A is defined to be $D\zeta/(S\zeta \cup D\zeta_{|B})$.

Proof: Previously we indexed the points in $T^*\Lambda_{r+1}N$ by $j \in \mathbb{Z}_r$, but for the purpose of this lemma, we index them by $0, 1, \ldots, r$ in $\mathbb{Z}$, and the points in $T^*\Lambda_r N$ by $0, 1, \ldots, r-1$. So we think of z_r as the extra point, and define the projection

$$\pi \colon T^*\Lambda_{r+1}N \to T^*\Lambda_r N$$

by forgetting z_r.

Recall the definition of A_{r+1}: we defined γ_j as the flow curve $\varphi_t(z_j), t \in [0, 1/(r+1)]$, but we could just as well have defined $\gamma_j = \varphi_t(z_j), t \in [0, t_j]$ where $\sum_j t_j = 1$ (and of course $z_j^- = \varphi_{t_j}(z_j)$). There is no difficulty in incorporating this in all the proofs so far if we assume something like $t_j < 2/r$ for all j. Also because $\sum_j t_j = 1$, the critical points will still be the 1-periodic orbits, but they are dissected differently. The critical values of these orbits also stay the same. This modified construction is needed because we wish to define an alternate A_{r+1} by $t_j = 1/r$ for $j \neq r$ and $t_r = 0$. Effectively this means that $z_0^- = z_r$. We can always change A_{r+1} back to the standard one by homotoping the t_j's, and get the same homotopy index because we have good index pairs during the homotopy for any interval (a, b).

Fixing all points $z_0, \ldots, z_{r-1}$ and q_r, but taking two different values of p_r, say p_{r1} and p_{r2}, we get

$$A_{r+1}(p_{r1}) - A_{r+1}(p_{r2}) = (p_{r1} - p_{r2})\varepsilon_{q_0} = (p_{r1} - p_{r2})\exp_{q_r}^{-1}(q_0), \qquad (9)$$

which implies that for all parameters except p_r fixed and $q_r = q_0$, the function A_{r+1} is constant, and the gradient $\nabla_{q_r} A_{r+1}$ as a function of p_r has a unique zero. We use this to define an embedding $i_r \colon T^*\Lambda_r N \to T^*\Lambda_{r+1}N$ by putting the new point $q_r = q_0$ and p_r equal to this unique critical point. Checking the terms in A_{r+1} and A_r we see that we have a commutative diagram

$$\begin{tikzcd}
T^*\Lambda_r N \arrow[rr, "i_r"] \arrow[dr, "A_r"'] & & T^*\Lambda_{r+1} N \arrow[dl, "A_{r+1}"] \\
& \mathbb{R} &
\end{tikzcd}$$

We defined i_r such that on the image we have $\nabla_{q_r} A_{r+1} = 0$, but from lemma 4.2 we see that also $\nabla_{p_r} A_{r+1} = 0$. Also, the embedding i_r maps critical points bijectively to critical points.

We would like to understand the Hessian with respect to variations of z_r on the embedding. This would give an understanding of how A_r behaves on a small normal bundle of the embedding. Equation (9) tells us that changing p_r changes the gradient $\nabla_{q_r} A_{r+1}$ by $-p_r$, and the expression in lemma 4.2 tells us that by changing q_r we approximately change ∇_{p_j} by $-q_r$, so in a normal chart and (q, p) splitting we approximately have the Hessian

$$\begin{bmatrix} 0 & -I \\ -I & 0 \end{bmatrix},$$

which is similar to

$$\begin{bmatrix} I & 0 \\ 0 & -I \end{bmatrix}.$$

So the negative eigenspace and positive eigenspace of this as a bundle over the embedding are both isomorphic to $\mathrm{Ev}_{q_0}^* TN$. This is a very good indication that the lemma is correct, but formally we need to define an index pair and a pseudo-gradient showing this. These can be constructed in much the same way as we constructed the index pair and pseudo-gradient in the proof of the above proposition. This case, however, is a bit more technical, because in the other case we had only negative directions in the normal bundle, but the essential ideas are the same. $\square$

We summarize the most important facts of this section in the following proposition.

Proposition 4.7. *There exist constants $K' > 0$ and $r_0 > 0$ such that: if H is any Hamiltonian with the properties and bounds described in the beginning of this section and $r > \max(K'(C_1 + C_2), r_0\mu)$ then*

$$I(A_r, X) = \mathrm{Th}(T\Lambda_r^\mu N).$$

Proof: We will prove that the K' from lemma 4.2 also works as K' in this case. Let H_0 be as in lemma 4.5, and choose $r_0 > C$ from that lemma such that we can use the result. Also, choose r_0 large enough for us to use lemma 4.3 on H_0. For this we need that $r_0 > K'(C^1_{H_0} + C^2_{H_0})$, but the bounds on H_0 can be chosen to be $C^1_{H_0} = \mu$ and $C_2 = C'\mu$, where C' is a constant depending only on the metric. So this choice is also linear in μ as the lemma requires. We define $H_t = (1-t)H_0 + tH$, and then by the assumption on r we can use lemma 4.3

on H_t for any t (we can use convex combinations of the bounds on H and H_0 as bounds on H_t). Thus if we chose a and b large enough to contain all the critical points of A^t_r for all $t \in I$, then by lemma 2.4 we see that the total index is defined for all t and is constant as a function of t. $\square$

5 Generalized Finite Reductions

In this section we introduce a more general family of finite reductions to the action integral.

Let $(M, \partial M)$ be a compact manifold with smooth boundary and with $M \subset M'$, where (M', ω) is an open symplectic manifold with $d\lambda = \omega$. Let g be a Riemannian structure on M' compatible with ω. We will work with a family of Hamiltonians $H_\delta \colon M' \to \mathbb{R}, 0 < \delta < \delta'$ which satisfy:

H1) There exist a neighborhood U of ∂M such that the Hamiltonian flow φ^δ_t of H_δ preserves U and is independent of δ.

H2) There is a constant C_1 which bounds $\|\nabla H_\delta\|$ for all δ, and the critical value associated to any 1-periodic orbit of H_δ is contained in the interval $(-\delta C_1, \delta C_1)$.

H3) There is a constant C_2 s.t. $\|\nabla^2 H_\delta\| < C_2 \delta^{-1}$.

Remark 5.1. This is somewhat technical. However, an important example to keep in mind of such a family is: for a closed manifold N with a riemannian structure and some $\mu > 0$ not a geodesic length and $\delta > 0$ small we define

$$H_\delta \colon T^*_{\varepsilon_1} N \to \mathbb{R},$$

where $T^*_{\varepsilon_1} N = \{(q,p) \in T^*N \mid \|p\| \le \varepsilon_1\}$ and $H(q,p) = h(\|p\|)$. We want $h(0) = h'(0) = 0$, and we want h'' to be a positive function with upper bound $2\mu\delta^{-1}$ and support in $[0, \delta]$, such that h' is constantly equal to μ on $[\delta, \varepsilon_1]$. Because any tangent to h intersects the y-axis above $-\mu\delta$ and below 0 the critical value of any 1-periodic orbit must lie in $[0, \delta\mu[$ (see remark 3.1). So H2 is fulfilled. For small δ we see that the flow is independent of δ on the set $\|p\| > \varepsilon_1/2$ and preserves it. Furthermore there are no 1 periodic orbits on this set so H1 is fulfilled. H3 is a simple calculation in the induced Riemannian structure on T^*N and is thus true in any Riemannian structure.

We want to define finite reductions of the action integral and prove existence of good index pairs for appropriate δs. The finite dimensional approximation of curves we will use in this section is

$$\Lambda_{r,\beta} M = \{\overrightarrow{z} \in M^r \mid e(\overrightarrow{z}) < \beta\},$$

where $e(\overrightarrow{z}) = e_g(\overrightarrow{z}) = r \sum_j \mathrm{dist}_g(z_j, z_{j+1})^2$ is the energy of the closed piecewise geodesic curve connecting the z_j's in cyclic order each piece parametrized by an interval of length $1/r$.

Definition 5.2. For any symplectic bundle $\xi \to M$ denote by $\mathcal{L}(\xi) \to M$ the fiber bundle with fiber $\mathcal{L}(\xi)_m$ the Grassmannian of Lagrangian subspaces of ξ_m. If ξ has a compatible inner product and the manifold has a Riemannian structure, we induce a Riemannian structue on $\mathcal{L}(\xi)$: each fiber is a Grassmannian of Lagrangian subspaces of a vector space with a metric, which means it has an induced Riemannian structure. We define the orthogonal complement to the fiber by parallel transport of the Lagrangian subspaces, and use the metric on M to define the inner product on this complement. The energy of a curve γ on this Riemannian manifold will by abuse of notation also be denoted $e(\gamma)$.

Our definition of the finite reduction will depend on a section $S_{\mathcal{L}}$ in the bundle

$$\Lambda\mathcal{L}(TM) \to \Lambda M. \tag{10}$$

Such a section may not exist so we now restrict to the case where it does. This section should have some bounds defined by a constant $C_{\mathcal{L}} > 0$: assume that $\gamma_s \in \Lambda M, s \in (-\varepsilon, \varepsilon)$ is a family of curves with energy less than β such that $\frac{\partial}{\partial s} e(\gamma_s)$ exists. We need $S_{\mathcal{L}}$ to fulfil

$$\begin{aligned} e(S_{\mathcal{L}}(\gamma_0)) &< C_{\mathcal{L}} \\ \sup_t \|\frac{\partial}{\partial s}(S_{\mathcal{L}}(\gamma_s))(t)_{|s=0}\| &< C_{\mathcal{L}} \sup_t \|\frac{\partial}{\partial s}(\gamma_s)(t)_{|s=0}\|, \end{aligned} \tag{11}$$

Note that the order in which we choose the data needed is: we first fix β then we choose a section $S_{\mathcal{L}}$ with such bounds depending on some $C_{\mathcal{L}}$. So we must make sure that when we choose β it is independent of $C_{\mathcal{L}}$.

Remark 5.3. In the case $M = T^*_{\varepsilon_1}N$ we have a canonical section in

$$\Lambda\mathcal{L}(T(T^*_{\varepsilon_1}N)) \to \Lambda T^*_{\varepsilon_1}N,$$

given by the vertical directions (the p-directions). This is the loop of a smooth section in $\mathcal{L}(T(T^*_{\varepsilon_1}N)) \to T^*_{\varepsilon_1}N$. So for any β it fulfils the bounds in (11).

Given a Lagrangian subspace $L \subset T_zM$ we define the *wedge* map

$$v_L \colon T_zM \to M$$

by decomposing $u \in T_zM$ as $u = u_L + u_{L^\perp}$ ($L^\perp = JL$) and then exponentiating u_L to a point in M say z'. Now parallel transport $u_{L^\perp}$ to z' using the exponential curve and exponentiate that to get z'' (see figure 3). We define $v_L(u) = z''$.



Figure 3: The wedge map.

This is a diffeomorphism at $u = 0$, so by compactness of M there is an $\varepsilon > 0$ such that: for any $z, z'' \in T^*_{\varepsilon_1}N$ with $\mathrm{dist}(z, z'') < \varepsilon$ and L a Lagrangian in

T_zM there is a unique $u \in T_zM$ close to 0 with $v_L(u) = z''$. In this case, we define the *L-curve* $\gamma^{\llcorner}(z, z'', L)$ by the curve starting at z'', going to z' by the exponential curve, and then continuing to z by the other exponential curve. We will later discuss parametrization. Notice that although z and z'' are in $M \subset M'$ we allow this L-curve to exit and reenter M.

We will use these L-curves to define our finite reductions. Figure 4 illustrates many aspects of the definition.



Figure 4: Curves involved in definition of finite reduction.

Definition 5.4. For any $\overrightarrow{z} \in \Lambda_{r,\beta}M$ let $S_{\mathcal{L}}(\overrightarrow{z})$ be the section evaluated at the piecewise geodesic connecting the z_j's in cyclic order. For large r we define

$$\begin{aligned} \gamma_j(t) &= \varphi_t^{\delta}(z_j), \quad t \in [0, 1/r] \\ z_j^- &= \gamma_{j-1}(1/r) \\ \gamma_j^{\llcorner} &= \gamma^{\llcorner}(z_j, z_j^-, S_{\mathcal{L}}(\overrightarrow{z})(j/r)) \end{aligned}$$

and

$$A_r(\overrightarrow{z}) = A_{r,g,S_{\mathcal{L}},\delta}(\overrightarrow{z}) = \sum_{j=1}^{r} \Big(\int_{\gamma_j} (\lambda - H_\delta dt) + \int_{\gamma_j^{\llcorner}} \lambda \Big).$$

We will also need

$$\begin{aligned} \varepsilon_{x_j} &= \exp_{z_j^-}^{-1}(z_j') \\ \varepsilon_{y_j} &= -\exp_{z_j}^{-1}(z_j') \\ \varepsilon_j &= \exp_{z_j^-}^{-1}(z_j) \end{aligned}$$

where z_j' is the point on $\gamma_j^{\llcorner}$ where the two geodesics meet. Finally we define an energy type functional

$$E(\overrightarrow{z}) = \sum_j \|\varepsilon_j\|^2$$

Notice that by using parallel transport it makes sence to write $\varepsilon_j \approx \varepsilon_{x_j} + \varepsilon_{y_j}$ for small ε_j.

We will prove existence of good pairs using the gradient of A_r and not a pseudo-gradient, and E will be used as a cut-off function. We have defined E such that it is zero if and only if the curves γ_j fits together to a 1-periodic orbit. Indeed E can be thought of as a finite version of the energy relative to the Hamiltonian flow:

$$E(\gamma) = \int_{s^1} \|\gamma'(t) - X_H(\gamma(t))\|^2 dt$$

In the case where $M = T^*_{\varepsilon_1}N$, $S_{\mathcal{L}} = S_{\mathcal{L}0}$ and the metric g is induced from a metric on N, we compare this to definition 4.1: the curve γ^{L}_j will be the curve first going in direction $\exp^{-1}_{q_j^-}(q_j)$, parallel transporting p_j^-, and after reaching the fiber over q_j it continues in a straight line in the fiber down to p_j. Integrating this over λ, one gets the term $p_j^- \varepsilon_{q_j}$. So this is indeed a generalization albeit only on a subset.

Remark 5.5. The function A_r can be approximated by $A_r \approx A_H \circ i_r$ where i_r is an embedding

$$i_r \colon \Lambda_{r,\beta}M \to \Lambda M.$$

We define i_r as the curve depicted in figure 4 with parametrization on the flow curves γ_j as defined, but the L-curves are parametrized on a much shorter interval - say $[0, s]$, where s is very small. Because s is very small we get almost no contribution from the integrating of the Hdt term over the L-curve and we approximately get the expression for A_r. This can be made more rigorous s.t. the homotopy indecies defined by such embeddings is the same as the one defined by A_r. This is, however, not needed but just a nice way to think about these functions, when relating them to Floer homology.

Lemma 5.6. *There exists a constant $K > 0$ independend of β and $C_{\mathcal{L}}$ (from equation* (11)*) and a constant $\delta_0 > 0$ such that: if $0 < \delta < \delta_0$ and $r > K\delta^{-1}$ then*

$$\nabla_{z_j} E = (\varepsilon_{x_{j+1}} - \varepsilon_{x_j}, \varepsilon_{y_{j+1}} - \varepsilon_{y_j}) + \frac{1}{10}\operatorname{bo}(\|\varepsilon_j\| + \|\varepsilon_{j+1}\|)$$

and

$$\nabla_{z_j} A_r = (\varepsilon_{y_j}, -\varepsilon_{x_{j+1}}) + \frac{1}{10}\operatorname{bo}(\|\varepsilon_j\| + \|\varepsilon_{j+1}\|) + r_j(\overrightarrow{z}),$$

where

$$\|\overrightarrow{r}(\overrightarrow{z})\| = \sqrt{\sum_j \|r_j(\overrightarrow{z})\|^2} \leq 2C_{\mathcal{L}}E.$$

Proof: We start by looking at $\nabla_{z_j} A_r$. If we move the curve $\overrightarrow{z}$ in some direction with unit speed all the Lagrangians $L_k = S_{\mathcal{L}}(\overrightarrow{z})(k/r) \subset T_{z_k}M$ changes with the points. We can ask how this movement is different from parallel transport at each point z_k. Because of the second bound in equation (11) we see that the rate of change of this difference is less than $C_{\mathcal{L}}$. In figure 5 we see that the change in symplectic area for small θ is bounded by



Figure 5: The change of symplectic area for small θ.

$$\sin(\theta)(\|\varepsilon_{x_j}\|^2 + \|\varepsilon_{y_j}\|^2) < 2\theta\|\varepsilon_j\|^2.$$

So the rate of change of A_r is less than

$$2C_{\mathcal{L}}\|\varepsilon_j\|^2$$

when moving the Lagrangian L_k with speed $C_{\mathcal{L}}$. So the contribution to the gradient of A_r from this has norm bounded by

$$2C_{\mathcal{L}}E$$

So the error in assuming that the change of L_k is given by parallel transport is no more than this. This is the error term denoted by $r_j(\overrightarrow{z})$ in the lemma.

So when calculating $\nabla_{z_j}A_r$ we can now assume that all the other Lagrangians $L_k, k \neq j$ are constant and L_j moves as parallel transport with respect to z_j.

We need a bound on E:

$$\begin{aligned} E(\overrightarrow{z}) = \sum_j (\operatorname{dist}(z_j, z_j^-))^2 \leq \sum_j (\frac{C_1}{r} + \operatorname{dist}(z_j, z_{j+1}))^2 < \\ < \sum_j \big(\tfrac{2(C_1)^2}{r^2} + 2\operatorname{dist}(z_j, z_{j+1})^2\big) < \frac{2C_1^2 + 2\beta}{r}. \end{aligned}$$

Because M is compact, we can choose ball shaped Darboux charts h for all points z such that $h(z) = 0$, and assume that they have a radius bounded from below by some small positive number. This is not a continuous family of charts, but just a choice for all points with this common bound. We can pick r large enough such that the points z_{j-1}, z_j^-, z_j , z_{j+1}^- and z_{j+1} all lie inside the chart centered at z_j. By composing the chart at z_j with a symplectic linear map we may assume that the differential $D_{z_j}h$ is an isometry. We can, by further composing with an element of $U(n)$, assume that the Lagrangian $L_j = h_* L_j \subset T_0\mathbb{R}^{2n}$ is equal to the imaginary part $i\mathbb{R}^n$ of $\mathbb{R}^{2n} \simeq \mathbb{C}^n$.

We now replace A_r with the function

$$f(z_j) = \int_{\gamma_{j-1}} (\lambda_0 - H dt) + \int_{\gamma_j^{\llcorner}} \lambda_0 + \int_{\gamma_j} (\lambda_0 - H dt) + \int_{\gamma_{j+1}^{\llcorner}} \lambda_0,$$

where we think of the points and curves as lying in $\mathbb{R}^{2n}$ and $\lambda_0 = \sum_j y_j dx_j$ is the standard 1-form in $\mathbb{R}^{2n}$. We assume that the metric is induced by the chart and the metric on M, so it is appriori not the standard one. The gradient of f at $z_j = 0$ is now the gradient $\nabla_{z_j}A_r$ which we wish to calculate.

As in the proof of lemma 4.2, we now replace f with f', which is f but dependend on the *independent* coordinates $z_j = (x_j, y_j)$ and $z_{j+1}^- = (x_{j+1}^-, y_{j+1}^-)$. Equation (4) used on the cotangent bundle $T^*\mathbb{R}^n \cong \mathbb{R}^{2n}$ calculates the gradient of the integral along γ_j (in the standard metric) to be

$$\nabla_{x_j, y_j, x_{j+1}^-, y_{j+1}^-}(\int_{\gamma_j} \lambda_0 - H dt) = (0, 0, y_{j+1}^-, 0). \tag{12}$$

First focus on the first two factors. because they are zero this is the gradient in any metric.

If the metric were flat, then ε_{x_j} and ε_{y_j} would both be linear and parallel to $\mathbb{R}^n$ and $i\mathbb{R}^n$ respectively, and we would get

$$\nabla_{x_j,y_j}\int_{\gamma_j^{\llcorner}}\lambda_0=\nabla_{z_j}\omega_0(\varepsilon_{x_j},\varepsilon_{y_j})=(\varepsilon_{y_j},0).$$

If the distance $\|\varepsilon_j\|=\operatorname{dist}(z_j^-,z_j)$ is very small this will approximate the linear case. So we get

$$\nabla_{x_j,y_j}\int_{\gamma_j^{\llcorner}}\lambda_0=(\varepsilon_{y_j},0)+\operatorname{bo}(c\|\varepsilon_j\|^2)$$

for some $c>0$. For large r we have $c\|\varepsilon_j\|<\frac{1}{10}$, and thus

$$\nabla_{x_j,y_j}\int_{\gamma_j^{\llcorner}}\lambda_0=(\varepsilon_{y_j},0)+\operatorname{bo}(\tfrac{1}{10}\|\varepsilon_j\|)$$

To calculate the other part of the gradient we linear translate the coordinate such that $z_{j+1}^-=0$. This changes the gradient of the individual integration parts, but not the overall gradient of f at the point z_{j+1}^-. Eventhough the metric at 0 for this new chart is not the standard metric it is C_1/r close to it, and so by similar arguments we see that for large r we get

$$\nabla_{z_{j+1}^-}\int_{\gamma_{j+1}^{\llcorner}}\lambda_0=(0,\varepsilon_{x_{j+1}})+\operatorname{bo}(\tfrac{1}{20}\|\varepsilon_{j+1}\|).$$

In these coordinates we also see that for the standard metric we have

$$\nabla_{z_j,z_{j+1}^-}(\int_{\gamma_j}\lambda_0-Hdt)=(-y_{j+1},0,0,0).\tag{13}$$

Again we use this to conclude that in our metric because the two last factors are also zero. Thus the actual gradient of f as a function of z_j is

$$\nabla_{z_j}f=(\varepsilon_{y_j},0)-\Phi_r(0,\varepsilon_{x_{j+1}})+\operatorname{bo}(\tfrac{1}{20}(2\|\varepsilon_j\|+\|\Phi_r^\delta\|\|\varepsilon_{j+1}\|))$$

where Φ_r^δ is the differential $D_{z_j}\varphi_{1/r}^\delta$. By increasing r this will be close to the identity in our coordinates. This is where we choose K, because if we change H, we also need to make r larger for Φ_r^δ to be close to the identity, and it is not difficult to prove that $r>K\delta^{-1}$ is good enough for some K depending only on the metric g and C_2 (from H3). So by making Φ_r^δ close to the identity and using that $\|\varepsilon_{x_{j+1}}\|<\|\varepsilon_{j+1}\|$ we get the wanted expression. All other bounds on r has not depended on δ, and can now be taken care of by decreasing δ_0.

The gradient $\nabla_{z_j}E$ is calculated similarly. □

Corollary 5.7. *Let $K>0$ be as in the previous lemma then there exists a $\delta_0>0$ such that: if $0<\delta<\delta_0$ and $r>K\delta^{-1}$ then*

$$\|\nabla E\|^2\leq 5E\leq 10\|\nabla A_r\|^2\leq 20E.$$

Equalities only if $E=0$.

This corollary has a very interesting implication: the critical points of A_r are the 1-periodic orbits regardless of the metric and $S_{\mathcal{L}}$. We will, however, later see that in the case of a non-degenerate critical point, the Morse index will depend on $S_{\mathcal{L}}$. In fact, the Morse index will turn out to be the Maslov index measuered with respect to $S_{\mathcal{L}}$ plus nr.

Proof: For any $\varepsilon > 0$ we can for large r (i.e. small δ_0) assume

$$(1-\varepsilon)\|\varepsilon_j\|^2 \leq (\|\varepsilon_{x_j}\|^2 + \|\varepsilon_{y_j}\|^2) \leq (1+\varepsilon)\|\varepsilon_j\|^2.$$

With $X_j = (\varepsilon_{x_{j+1}} - \varepsilon_{x_j}, \varepsilon_{y_{j+1}} - \varepsilon_{y_j})$ we see that

$$\begin{aligned} \|\nabla E\|^2 &\leq \sum_j (\|X_j\|^2 + \tfrac{1}{10}\|X_j\|(\|\varepsilon_j\| + \|\varepsilon_{j+1}\|) \leq \\ &\leq (1+\varepsilon)4E + \tfrac{\sqrt{1+\varepsilon}8E}{10} \leq 5E. \end{aligned}$$

With $X_j = (\varepsilon_{y_j}, -\varepsilon_{x_{j+1}})$ we see that

$$\begin{aligned} \|\nabla A_r\|^2 &\leq \sum_j \|X_j\|^2 + \tfrac{1}{10}\|X_j\|(\|\varepsilon_j\| + \|\varepsilon_{j+1}\|) = \\ &\leq (1+\varepsilon)E + \tfrac{\sqrt{1+\varepsilon}}{5}E \leq 2E. \end{aligned}$$

and

$$\begin{aligned} \|\nabla A_r\|^2 &\geq \sum_j \|X_j\|^2 - \tfrac{1}{10}\|X_j\|(\|\varepsilon_j\| + \|\varepsilon_{j+1}\|) = \\ &\geq (1-\varepsilon)E - \tfrac{\sqrt{1+\varepsilon}}{5}E \geq \tfrac{E}{2}. \end{aligned}$$

□

We will also need cut-off functions keeping the z_j's away from the boundary of M. So we define

$$Q(z) = -\operatorname{dist}(z, \partial M), \qquad z \in M.$$

This is not smooth on all of M. However, we only need it to be smooth near the boundary. Because M is compact we can find $\tau > 0$ such that Q is smooth on the colar

$$B_\tau = \{z \in M \mid Q(z) \geq -\tau\}.$$

By possibly making τ smaller we may also assume that $B_\tau \subset U$, where U is the set on which H_δ has no 1-periodic orbits and is independent of δ. We then define the functions $Q_j(\overrightarrow{z}) = Q(z_j)$. The next lemma ensures that we can use these as cut-off functions.

Lemma 5.8. *Let K be as in lemma 5.6. There exists $k > 0$ and $\delta_0 > 0$ such that: If $0 < \delta < \delta_0$ and $r > K\delta^{-1}$, then for any $\overrightarrow{z} \in \Lambda_{r,\beta}M$ with $z_j \in B_\tau$ we have*

$$|\nabla A_r \cdot \nabla Q_j| < \sqrt{r}k\|\nabla A_r\|^2.$$

Proof: We see that $\|\nabla Q_j\| = \|\nabla_{z_j} Q_j\| = 1$ when $z_j \in B_\tau$. If we have a lower bound of the type

$$\|\nabla A_r\| > \frac{c}{\sqrt{r}}, \tag{14}$$

on the same set then

$$|\nabla A_r \cdot \nabla Q_j| \leq \|\nabla A_r\| < \sqrt{r} c^{-1} \|\nabla A_r\|^2.$$

So we need such a bound. If we pick δ_0 smaller than the above corollary needs then $2\|\nabla A_r\|^2 \geq E$. So a lower bound on E of the type c/r will do the trick. This follows if we show that $s = \sum_j \|\varepsilon_j\| > \sqrt{c}$.

Define the function $L_H \colon \Lambda M \to \mathbb{R}$ by

$$L_H(\gamma) = \int_\gamma \|\gamma' - X_H\| dt.$$

This is zero if and only if γ is a 1-periodic orbit of the Hamiltonian flow. We can approximate s by using L_H in the following way: let γ be the closed curve which is the flow curve $\varphi_{t-j/r}(z_j)$ when $t \in [j/r, (j+1-1/k)/r]$ and when $t \in [(j+1-1/k)/r, (j+1)/r]$ it is the geodesic connecting $\varphi_{(1-1/k)/r}(z_j)$ and z_{j+1}. As k tends to infinity $L_H(\gamma)$ tends to s. So all we need is a lower bound on L_H.

This is obtained by relating it to a more well-known functional. Let $\gamma_2(t) = \varphi^\delta_{-t}(\gamma(t))$. We now divide into two cases. First case: if γ_2 is a curve in U then

$$L_H(\gamma) > C \int_0^1 \|D_{\gamma(t)} \varphi^\delta_{-t}(\gamma'(t) - X_{\gamma(t)})\| dt = C \int_0^1 \|\gamma_2'(t)\| dt > c$$

where $C < \|D_z \varphi^\delta_t\|$ for all $z \in U$, which is independent of δ by assumption H1. The last inequality is because H1 implies that $C \operatorname{dist}(\gamma_2(0), \gamma_2(1)) > c$ also independent of δ. If γ_2 is not totally contained in U then the above inequality only works on the pieces of γ_2 that is contained in U, but then we see that γ_2 has to exit U and reenter. So integrating over the part of γ_2 which is in U we get the bound $L_H(\gamma) > C(\operatorname{dist}(\gamma_2(0), M - U) + \operatorname{dist}(M - U, \gamma_2(1))) > c$. $\square$

Proposition 5.9. *There exists constants* $K >> 0, \beta >> 0$ *such that: for any section* $S_\mathcal{L}$ *in the bundle in* (10) *satisfying the bounds* (11) *there exists a* $\delta_0 > 0$ *so that the following holds.*

For any $0 < \delta < \delta_0$ *and* $r \in [K\delta^{-1}, 2K\delta^{-1}]$ *we have a good index pair for the total index of* $A_r \colon \Lambda_{r,\beta} M \to \mathbb{R}$*, and for all small* C^1*-perturbations of* H*, the good index pair will contain all of* A_r*'s critical points.*

Proof: We will use the formula in lemma 2.5 to create good index pairs using E and the functions Q_j as cut-off functions. We need these functions to seperate the pair we construct from the boundary of $\Lambda_{r,\beta} M$. That is, we want a good index pair in the interior of $\Lambda_{r,\beta} M$.

First Choose K as large as needed in corollary 5.7. Then choose $\beta = 32 C_1 K + 2C_1^2$ and $\delta_0 > 0$ as small as the corollary needs for this K, β and $C_\mathcal{L}$.

The corollary tells us that that $\nabla A_r \cdot \nabla E < \sqrt{10} \nabla A_r \cdot \nabla A_r$, and that $E = 0$ for critical points.

The assumption H2 tells us that the critical values of A_r is bounded in the interval $(-\delta C_1, \delta C_1)$. So we choose $-a = b = \delta C_1$, such that the total index of A_r is the same as $I_a^b(A_r)$. We will use E as a cut-off function with $0 < s$ and $t - s >= \sqrt{10}(b - a)$. This means we can choose t_j less than $4(b - a) = 8\delta C_1 \leq 16 C_1 K/r$. Knowing that $E < 16 C_1 K/r$ will provide a bound on the standard energy:

$$e(\overrightarrow{z}) = r \sum_j \operatorname{dist}(z_j, z_{j+1})^2 < r \sum_j (2 \operatorname{dist}(z_j, z_j^-)^2 + \tfrac{2\mathbb{C}_1^2}{r^2}) <$$
$$< 2rE + 2C_1^2 < 32 C_1 K + 2C_1^2 < \beta.$$

We now see that using E as a cut-off function will keep the pair seperated from the part of the boundary of $\Lambda_{r,\beta} M$ given by $e(\overrightarrow{z}) = \beta$.

To keep us away from the boundary of $\Lambda_{r,\beta} M$ with $z_j \in \partial M$ we use the functions Q_j from the previous lemma. We define $s_j = 2\tau/3$ and $t_j = \tau/3$. Looking at lemma 2.5 again, we see that we need

$$\nabla A_j \cdot \nabla Q_j < \tfrac{t_j - s_j}{b - a} \|\nabla A_r\|^2 = \tfrac{\tau}{6 C_1 \delta} \|\nabla A_r\|^2.$$

This is easily achived for small $\delta > 0$ using the previous lemma and the fact that $\delta^{-1} > r/(2K)$.

The last statement in the lemma follows from the fact that E is continuously dependent on $H \in C^1(M', \mathbb{R})$ and for critical points we know $E = 0$. So for a small enough C^1 pertubation we can assume that the newly formed critical points of A_r are close to the old ones. $\square$

6 Stabilization of Generalized Finite Reductions

Let $(M, \partial M) \subset M'$ and $H_\delta^M, \delta > 0$ fulfil H1 through H3 from the previous section. Define $P = M \times (D^{2k}, \omega_0)$ and $P' = M' \times \mathbb{R}^{2k}$ for some k and ω_0 the standard symplectic form on $\mathbb{R}^{2k}$. Also define $H_\delta \colon P' \to \mathbb{R}$ by $H_\delta(z_1, z_2) = H_\delta^M(z_1) + H_D(z_2)$, where

$$H_D(z_2) = \|z_2\|^2.$$

The Hamiltonian flow for H_D is circular around 0 with revolution time 2π, but we only flow for a time period of 1, so the only orbit is 0, and this orbit has action 0. So the 1-periodic orbits for H_δ are the 1-periodic orbits for H_δ^M on the first factor and constantly equal to 0 on the second factor, and the critical value of these orbits are the same as on M'. We conclude that $(P, \partial P)$, P' and H_δ also fulfil H1 through H3 from the previous section. The only difference is that P have co-dimension 2 boundary components, and in the previous section we assumed the boundary of M to be smooth.

For us to define finite reductions as in the previous section we still require a section

$$S_{\mathcal{L}} \colon \Lambda(M \times D^{2k}) \to \Lambda\mathcal{L}(TM \times (R^{2k}, \omega_0))$$

with the same type of bounds (11) as before on curves with energy less than β. We also require a compatible metric g on the space P'. However, in this section we assume that the metric is the product of a compatible metric with the standard metric on $\mathbb{R}^{2k}$. The corresponding finite version of the loop space $\Lambda_{r,\beta}(M \times D^{2k})$ will consist of curves denoted by $\overrightarrow{z} = (\overrightarrow{z_1}, \overrightarrow{z_2})$. So that $\overrightarrow{z_1}$ consists of points in M and $\overrightarrow{z_2}$ in D^{2k}.

Definition 6.1. A section $S_{\mathcal{L}}$ is said to be of *product type* if it factors through

$$\Lambda M \quad \text{and} \quad \Lambda(\mathcal{L}(TM) \times \mathcal{L}(k)),$$

where $\mathcal{L}(k)$ is the Grassmannian of Lagrangian subspaces in $\mathbb{R}^{2k}$.

Factoring through the first of the two spaces is equivalent to the section not depending on the second (contractible) factor ΛD^{2k}. Factoring through the second space is equivalent to all Lagrangians defined by the section split as direct sums of two Lagrangians, one in each factor. On the level of fibers this corresponds to the section being in the subset defined by $(\Lambda\mathcal{L}(n)) \times (\Lambda\mathcal{L}(k)) \subset \Lambda\mathcal{L}(n+k)$.

For the rest of this section, $S_{\mathcal{L}}$ will be of product type and A_r will be the finite reduction defined as in the previous section on $\Lambda_{r,\beta}P$. In this case A_r splits into two factors

$$A_r(\overrightarrow{z_1}, \overrightarrow{z_2}) = A_r^M(\overrightarrow{z_1}) + A_r^D(\overrightarrow{z_1}, \overrightarrow{z_2})$$

where A_r^M is the function defined in the previous section on $\Lambda_{r,\beta}M$ by restricting our section to the first factor, and $A_r^D(\overrightarrow{z_1}, -)$ is the finite reduction on D^{2k} defined by the the Hamiltonian H_D and the constant section: the restriction of $S_{\mathcal{L}}(\overrightarrow{z_1})$ to the second factor.

This gives us

$$\begin{aligned} \nabla_{\overrightarrow{z_1}} A_r &= \nabla A_r^M + \nabla_{\overrightarrow{z_1}} A_r^D \\ \nabla_{\overrightarrow{z_2}} A_r &= \nabla_{\overrightarrow{z_2}} A_r^D \end{aligned} \tag{15}$$

It would be nicer if we did not have the second term in the first line, and looking at the proof of lemma 5.6, one sees that this is part of the error term, so some how it should not be relevant $\overrightarrow{r}(\overrightarrow{z})$. We will in fact show that

$$X = X_1 \oplus X_2 = \nabla_{\overrightarrow{z_1}} A_r^M \oplus \nabla_{\overrightarrow{z_2}} A_r^D,$$

is a pseudo-gradient and as in proposition 5.9 we have good index pairs. This is made more precise and more general in the following technical lemma, which is a small generalization of proposition 5.9 to this product case.

Lemma 6.2. *There exists constants $K >> 0, \beta >> 0$ such that: for any section $S_{\mathcal{L}}$ in the bundle in* (10) *satisfying the bounds* (11) *there exists a $\delta_0 > 0$ so that the following holds.*

For any $0 < \delta < \delta_0$, $r \in [K\delta^{-1}, 2K\delta^{-1}]$, $0 < s \leq 1$ and $0 \leq t \leq 1$ we have a good index pair for the total index $I(A_r, X)$ of $A_r \colon \Lambda_{r,\beta}M \to \mathbb{R}$, where

$$X = (s\nabla A_r^M + t\nabla_{\overrightarrow{z_1}} A_r^D) \oplus \nabla_{\overrightarrow{z_2}} A_r^D$$

Proof: First we notice that the energy splits

$$E(\overrightarrow{z}) = E_1(\overrightarrow{z}) + E_2(\overrightarrow{z}),$$

where $E_1(\overrightarrow{z}) = E(\overrightarrow{z_1}, 0)$ and $E_2(\overrightarrow{z}) = E(\{*\}, \overrightarrow{z_2})$. So the gradient of E is

$$\nabla E = \nabla E_1 \oplus \nabla E_2.$$

Because of the bounds in equation (11), taking a variation of $\overrightarrow{z_1}$ in a unit direction will not change the angles of the Lagrangians on the second factor by more than $C_{\mathcal{L}}$. This changes A_r^D by no more than $C_{\mathcal{L}} E_2$. Indeed, this is the same argument as in the proof of lemma 5.6 and is explained in figure 5. So the norm of the gradient $\nabla_{\overrightarrow{z_1}} A_r^D$ is less than $C_{\mathcal{L}} E_2$. Using this, equation (15) and corollary 5.7 on both E_1, A_r^M and $E_2, A_r^D(\overrightarrow{z_1}, -)$ we get

$$\begin{aligned} X \cdot \nabla E &\leq \big(s\|\nabla_{\overrightarrow{z_1}} A_r^M\| + t\|\nabla_{\overrightarrow{z_1}} A_r^D\|\big)\|\nabla E_1\| + \|\nabla_{\overrightarrow{z_2}} A_r^D\|\|\nabla E_2\| \\ &\leq \big(s\sqrt{2E_1} + tC_{\mathcal{L}} E_2\big)\sqrt{5E_1} + \sqrt{2E_2}\sqrt{5E_2} \\ &\leq 4(sE_1 + tC_{\mathcal{L}} E_2\sqrt{E_1} + E_2) \end{aligned}$$

and

$$\begin{aligned} X \cdot \nabla A_r \quad &\geq (s\nabla_{\overrightarrow{z_1}} A_r^M + t\nabla_{\overrightarrow{z_1}} A_r^D) \cdot (\nabla_{\overrightarrow{z_1}} A_r^M + \nabla_{\overrightarrow{z_1}} A_r^D) + \|\nabla_{\overrightarrow{z_2}} A_r^D\|^2 \\ &\geq sE_1/2 - (t+s)\sqrt{E_1/2}C_{\mathcal{L}} E_2 - t(C_{\mathcal{L}} E_2)^2 + E_2/2. \end{aligned} \tag{16}$$

By making δ_0 smaller we make r larger and this makes $E = E_1 + E_2$ smaller. So for small δ_0 we can assume

$$X \cdot \nabla E \leq 5(sE_1 + E_2) \leq 20 X \cdot \nabla A_r.$$

This proves that X is a pseudo-gradient, because at non-critical points we have $E_1 + E_2 > 0$. It also proves that we can use E as a cut-off function in the same way we did in the proof of proposition 5.9.

Similarly, we define functions Q_j^M to be the distance of $\overrightarrow{z_1}$ to the boundary ∂M, and so by using lemma 5.8 on A_r^M and Q_j^M we get on the set $Q_j^M(\overrightarrow{z}) > -\tau$ that

$$\begin{aligned} X \cdot \nabla Q_j^M &< s\nabla_{\overrightarrow{z_1}} A_r^M \cdot \nabla Q_j^M + tC_{\mathcal{L}} E_2\|\nabla Q_j^M\| \\ &< s\sqrt{r}k\|\nabla_{\overrightarrow{z_1}} A_r^M\|^2 + 2tC_{\mathcal{L}} E_2 c \\ &< \sqrt{r}k(X \cdot \nabla A_r). \end{aligned}$$

To get the last inequality we may have to further increase r.

Similarly we define the function Q_j^D to be the distance of $\overrightarrow{z_2}$ to the boundary of D^{2k} and use lemma 5.8 to get

$$X \cdot \nabla Q_j^D < \sqrt{r}kX \cdot \nabla_{\overrightarrow{z_2}} A_r^D$$

Now we have enough cut-off functions to get a compact pair in the interior of $\Lambda_{r,\beta}(N \times D^{2k})$, just as in the proof of proposition 5.9. $\square$

Remark 6.3. Because $\nabla_{uz_2} H_D = u\nabla_{z_2} H_D$, we see that flow curves for the Hamiltonian flow of H_D is preserved under scaling. So if γ is a flow curve then $u\gamma$ is a flow curve. This means that the curve over which we integrate λ_0 scales proportionally with $\overrightarrow{z_2}$, so the integral of the canonical 1-form scales quadratically. Furthermore, H_k is quadratic. So we conclude

$$A_r^D(\overrightarrow{z_1}, u\overrightarrow{z_2}) = u^2 A_r^D(\overrightarrow{z_1}, \overrightarrow{z_2}). \tag{17}$$

Because this is a smooth function it must be equal to its Hessian at 0. So $A_r^D(\overrightarrow{z_1}, -)$ is in fact a quadratic form in $\overrightarrow{z_2}$.

If the critical point 0 were degenerate it would not be an isolated critical point. So it is in fact non-degenerate.

Definition 6.4. We define $\zeta \to \Lambda_{r,\beta} M$ to be the vector bundle with fiber at $\overrightarrow{z_1}$ the negative eigenspaces for the quadratic form $A_r^D(\overrightarrow{z_1}, -)$.

Lemma 6.5. *Assume that $K, \beta, \delta_0, \delta$ and r satisfy the conditions in the previous lemma. Then the total homotopy index of A_r is the relative Thom suspension of ζ of the homotopy index of A_r^M.*

Note that the relative Thom suspension of A/B of a bundle ζ over A is defined to be $D\zeta/(S\zeta \cup D\zeta_{|B})$.

Proof: First choose an index pair (A, B) for A_r^M, with $a = -C_1\delta$ and $b = C_1\delta$ (cf H2 in the previous section). We will extend this to an index pair for A_r. Let $E^{\pm}_{\overrightarrow{z_1}}$ be the negative/positive eigenbundle of $A_r^D(\overrightarrow{z_1}, -)$. It is easy to construct index pairs very close to zero for a non-degenerate quadratic form on D^{2kr}, so we do this fiber-wise

$$A_{\overrightarrow{z_1}} = D_\varepsilon E^-_{\overrightarrow{z_1}} \times D_\varepsilon E^+_{\overrightarrow{z_1}}$$
$$B_{\overrightarrow{z_1}} = S_\varepsilon E^-_{\overrightarrow{z_1}} \times D_\varepsilon E^+_{\overrightarrow{z_1}}.$$

Since $e(A) \in [0, \beta)$ and A is compact we have $e(A) \in [0, \beta - c]$, so we can find an $\varepsilon > 0$ such that $A_{\overrightarrow{z_1}}$ is contained in $\Lambda_{r,\beta}(N \times D^{2k})$ for all $\overrightarrow{z_1} \in A$. Similarly we can assume for small $\varepsilon > 0$ that A_r on these sets assumes only values in $(-2C_1\delta, 2C_1\delta)$. Define

$$A' = \bigcup_{\overrightarrow{z_1} \in A} A_{\overrightarrow{z_1}}$$
$$B' = (\bigcup_{\overrightarrow{z_1} \in B} A_{\overrightarrow{z_1}}) \cup (\bigcup_{\overrightarrow{z_1} \in A} B_{\overrightarrow{z_1}})$$

for such an ε.

Claim: We can find $0 < s \leq 1$ and put $t = 0$ in the previous lemma such that (A_r, X) has this as an index pair. I1 and I2 from the definition of index pair have been taken care of. I3 is because critical points of A_r are of the form $(\overrightarrow{z_1}, 0)$, where $\overrightarrow{z_1}$ is a critical point for A_r^M. To get I4 we use s, because s controls the speed of the flow on the first factor. For s very small, any point in B' will on the second factor flow entirely out of $\Lambda_{r,\beta}(M \times D^{2k})$ before the quadratic form $A_r(\overrightarrow{z_1}, -)$ changes too much. Because B' is compact we can choose s such that this is true for all $b \in B'$.

The quotient A'/B' is the wanted homotopy index. However, the statement in the lemma is for any t and s, but this follows because the previous lemma makes sure that the homotopy index is independent of s and t. $\square$

For us to use this lemma we would like to be able to compute the isomorphism class of the negative eigenbundle, and the next lemma helps us do just that.

Lemma 6.6. *Let $B_r^D(\overrightarrow{z_1},-)$ be the quadratic form defined like $A_r^D(\overrightarrow{z_1},-)$, but with the Hamiltonian $H_D=0$. Let $\zeta_B \to \Lambda_{r,\beta}M$ be the bundle with fiber the negative eigen bundle of $B_r^D(\overrightarrow{z},-)$ then*

$$\zeta \cong \zeta_B \oplus \mathbb{R}^{2k}.$$

Note that B_r^D is degenerate in each fiber, because in the fiber over $\overrightarrow{z_1}$, the quadratic form $B_r^D(\overrightarrow{z_1},-)$ is a finite reduction, which has the $2k$ dimensional subspace consisting of the constant curves as degenerate points (the 1-periodic orbits). We did not use this Hamiltonian in the definition of our finite reduction because we wanted index pairs, but the negative eigenbundle is easier to calculate for the quadratic form B_r^D.

Proof: We define a continuous family $A^s(\overrightarrow{z_1},-)$ of quadratic forms on $\Lambda_{r,\beta}D^{2k}$ by finite dimensional reduction: we use the Hamiltonians $H_s=s\|z_2\|^2$ and the last factor of the section $S_{\mathcal{L}}(\overrightarrow{z_1})$. Then $B_r^D(\overrightarrow{z_1},-)=A^0(\overrightarrow{z_1},-)$ and $A_r^D(\overrightarrow{z_1},-)=A^1(\overrightarrow{z_1},-)$. The argument in remark 6.3 proves that these are quadratic forms. So from now on we consider only their Hessians defined as quadratic forms on $\mathbb{R}^{2nr}$. Since the Hamiltonian flow for H_s has the trivial 1-periodic orbit 0 for $0<s<2\pi$, these are non-degenerate. So they have isomorphic negative eigenbundles, but we need to see what happens at $s=0$. The critical points of A^0 are precisely the constant curves, so the Hessian is degenerate, and the kernel as a bundle over $\Lambda_{r,\beta}T^*N$ is the trivial bundle of dimension $2k$. We prove the lemma by proving that for a small perturbation of $s=0$ in positive direction, this kernel becomes part of the negative eigenspace. This is a point-wise calculation, so assume $\overrightarrow{z_1}$ is fixed.

Denote by E_-, E_0 and E_+ the negative, zero and positive eigenspace of A^0. It is enough to prove that the first order change in s at $s=0$ of A_r^s is negative definite on the kernel E_0. Indeed, if so we can restrict A_r^0 to the sphere of $E_0\oplus E_-$ and what we see is a non-positive function on a closed manifold, which is then perturbed to the first order to be negative on the set where it is zero. This will imply that the function is in fact going to be negative on the entire sphere for very small s, and thus A_r^s is negative definite on $E_0\oplus E_-$ for small $s>0$.

So to prove this negativity on E_0, we look at A_r^s on E_0 for s close to zero. The kernel E_0 is the set of constant curves, so we assume that $z_j=z_{j+1}$ for all $j\in\mathbb{Z}_r$. We need to take a look at the precise definition of

$$A^s=\sum_j\Big(\int_{\gamma_j}\lambda_0-H_s dt+\int_{\gamma_j^{\llcorner}}\lambda_0\Big)$$

For $s=0$ all of this is zero (on E_0) because γ_j and $\gamma_j^{\llcorner}$ are constant, and H_s is zero. We want to prove that the dominating term when perturbing to positive s is $-H_s$, which is negative.

Because any time independent Hamiltonian is constant on its flow curves, we can rewrite this as

$$\begin{aligned} A^s &= \int_{\sum_j(\gamma_j+\gamma_j^{\llcorner})} \lambda_0 + \frac{1}{r}\sum_j H_s(z_j) = \\ &= \int_{\sum_j(\gamma_j+\gamma_j^{\llcorner})} \lambda_0 - H_s(z_0). \end{aligned}$$

The curves γ_j are the $1/r$ time flow curves of H_s, so they have lengths of order $\|\nabla H_s\|/r$ which is of order $s\|z_0\|/r$, and since $z_j = z_{j+1}$, and $\gamma_j^{\llcorner}$ connects the endpoint of γ_j with z_{j+1}, the same is true for $\gamma_j^{\llcorner}$. This means that the integral, which is the symplectic area enclosed by the closed curve obtained by concatenating γ_j and $\gamma_j^{\llcorner}$, is of order $(s\|z_0\|/r)^2$. We have r of these terms summed, but this is still of order $(s\|z_0\|)^2/r$. The term $H_s(z_0)$ is equal to $s\|z_0\|^2$, so this is the dominating term in s and the lemma follows. $\square$

7 The Maslov Bundle and Index of Finite Reductions

Let $j\colon L \to T^*N$ be a Lagrangian embedding. We will define the Maslov bundle relative to this embedding. It is a generalization of the Maslov index related to curves of Lagrangian subspaces in $\mathbb{R}^{2n}$ (see e.g. [7]). In fact the bundle is a canonical virtual vector bundle over ΛL, such that the dimension of this bundle on each component is precisely the Maslov index.

The projection $T^*N \to N$ will be denoted π. For any point $q \in L$ the tangent space T_qL is mapped by j_* to a Lagrangian subspace of $T_{j(q)}(T^*N)$, and by abuse of notation we define

$$j_*\colon L \to \mathcal{L}(T(T^*N)),$$

where $\mathcal{L}(T(T^*N))$ is defined in definition 5.2. A stabilization of this map with a vector bundle $\zeta \to N$ will be denoted by $j_* \oplus \zeta$ and is defined by

$$(j_* \oplus \zeta)(q) = j_*(T_qL) \oplus (\pi^*\zeta) \subset T_{j(q)}(T^*N) \oplus (\pi^*\zeta) \oplus (\pi^*\zeta)^*,$$

which is also Lagrangian (in the obvious symplectic structure).

If ζ is the normal bundle of N for some embedding $i_N\colon N \to \mathbb{R}^n$, we get a canonical symplectic trivialization

$$\eta_i\colon T(T^*N) \oplus (\pi^*\zeta) \oplus (\pi^*\zeta)^* \to T^*N \times (\mathbb{R}^{2n}, \omega_0).$$

This is defined by using the Riemannian metric induced from i_N to split the tangent space of T^*N at z into $T_{\pi(z)}N \oplus T^*_{\pi(z)}N$, then mapping $V_z = T_{\pi(z)}N \oplus \zeta_{\pi(z)}$ isomorphically to $\mathbb{R}^n$ by the obvious map, and mapping $V_z^* = T^*_{\pi(z)}N \oplus \zeta^*_{\pi(z)}$, by the inverse of the dual to this map, to $i\mathbb{R}^n$. If we compose this trivialization with $j_* \oplus \zeta$ we get a map from L to $\mathcal{L}(n)$ (the Grassmannian of Lagrangian subspaces in $\mathbb{R}^{2n}$), and since all embeddings are isotopic for n sufficiently large, we have a map unique up to homotopy

$$J_*\colon L \to \mathcal{L}(n).$$

If we stabilize this map to get a map to $\mathcal{L}(2n)$, then we can homotope it to be a map into $\mathcal{L}(n) \times \mathcal{L}(n) \subset \mathcal{L}(2n)$ which is constant on the first factor. This means that if we further stabilize this map by the tangent space of T^*N and choose to trivialize this copy of T^*N with its normal bundle, we have a different interpretation of the map J_*: For any point $l \in L$ we have two different Lagrangian subspaces of $T_l(T^*N)$, the tangent space of L and the horizontal part of $T(T^*N)$. So we have two different sections in the bundle $\mathcal{L}(T(T^*N))_{|L}$. If we stabilize both sections with $\mathbb{R}^n \subset \mathbb{R}^{2n}$ to make them bundles of Lagrangian subspaces in a higher dimensional symplectic bundle over L, we can - if n is large enough - homotope their "difference" through Lagrangian subspaces to the $\mathbb{R}^{2n}$ component. It is this difference that J_* measures.

Because $\lim_{n\to\infty} \mathcal{L}(n) = \mathcal{L} \simeq \mathrm{U}/\mathrm{O} \simeq \Omega^6\mathrm{O}$ (see e.g. [7] for the first homotopy equivalence and [8] for the last) is an H-space we have a homotopy equivalence

$$\mathrm{Ev}_0 \times \pi_\Omega \colon \Lambda\mathcal{L} \to \mathcal{L} \times \Omega\mathcal{L}, \tag{18}$$

where Ev_0 is evaluation at the base point, and π_Ω is homotopic to point-wise multiplication with the homotopy inverse of Ev_0.

Definition 7.1. The Maslov bundle η is the virtual bundle defined by the map

$$\Lambda L \xrightarrow{\Lambda J_*} \Lambda\mathcal{L}(n) \longrightarrow \Lambda\mathcal{L} \xrightarrow{\pi_\Omega} \Omega\mathcal{L} \xrightarrow{\simeq} \mathbb{Z} \times B\mathrm{O}.$$

The latter equivalence comes from Bott periodicity (see e.g. [8]). The $\mathbb{Z}$ corresponds to the dimension of the virtual bundle or $\pi_1(\mathcal{L})$, the latter being one of the many definitions of the Maslov index (see [7]).

We want to use this to calculate the type of homotopy indices defined in the previous section. To do this we need to put the loop of J_* on a standard form.

For any $V_+ \subset \mathbb{R}^n$, $V_- \subset \mathbb{R}^n$ and $V_0 \subset \mathbb{R}^n$ pairwise orthogonal and $V_+ \oplus V_- \oplus V_0 = \mathbb{R}^n \subset \mathbb{C}^n$, we define the curve $\gamma_{(V_+,V_-,V_0)} \in \Omega\mathcal{L}(n)$ by

$$\gamma_{(V_+,V_-,V_0)}(t) = e^{i\pi t}V_+ \oplus e^{-i\pi t}V_- \oplus V_0 \in \mathcal{L}(n),$$

for $t \in [0,1]$. The space of such curves will be denoted $\Omega_s\mathcal{L}(n)$ (s for standard form). Over this space we have the three canonical bundles V_+, V_- and V_0.

Lemma 7.2. *Any map $f \colon K \to \Lambda\mathcal{L}$, where K is of compact homotopy type, can be homotoped to factor through*

$$\mathcal{L}(n_1) \times \Omega_s\mathcal{L}(n_2) \subset \Lambda\mathcal{L}(n_1+n_2) \to \Lambda\mathcal{L}$$

for large enough n_1 and n_2. Furthermore, the virtual bundle defined by the map f over K by composing it with π_Ω is the pullback of $V_+ - V_-$.

Proof: The map factors through $\mathcal{L}(n_1) \times \Omega\mathcal{L}(n)$ because the map in equation (18) is a homotopy equivalence and because K is of compact homotopy type. So we can consider a map f' to $\Omega\mathcal{L}(n)$ and show that it factors through $\Omega_s\mathcal{L}(n)$ for large n. We may have to stabilize by a number of Ωi_n's, where

$$i_n \colon \mathcal{L}(n) \to \mathcal{L}(n+1)$$

is the standard stabilization which gives the limit $\mathcal{L}$.

This part of the proof is standard Morse theory as in the proof of Bott periodicity (see e.g. [8]): multiplication with $e^{-i\pi t/2}$ on the curves in $\mathcal{L}(n)$ gives a homeomorphism of $\Omega\mathcal{L}(n) = \Omega(\mathcal{L}(n), \mathbb{R}^n, \mathbb{R}^n)$ to $\Omega(\mathcal{L}(n), \mathbb{R}^n, i\mathbb{R}^n)$. We now think of f' as mapping into this space. In [8] part IV paragraph 24 the space of minimal geodesics for this space is computed to be (with some interpretation into the current context)

$$\Omega_m(n) = \Omega_m(\mathcal{L}(n), \mathbb{R}^n, i\mathbb{R}^n) = \{\gamma \mid \gamma(t) = e^{i\pi t/2}W \oplus e^{-i\pi t/2}W^\perp, W \subset \mathbb{R}^n\}.$$

The embedding of this space into $\Omega(\mathcal{L}(n), \mathbb{R}^n, i\mathbb{R}^n)$ has high connectivity on the components where $\dim(W)$ and $\dim(W^\perp)$ are both high.

To get high connectivity we stabilize f' by

$$\gamma(t) = e^{i\pi t/2}\mathbb{R} \oplus e^{-i\pi t/2}\mathbb{R} \subset \mathbb{R}^4.$$

That is, we compose with the map

$$\oplus\gamma \colon \Omega\mathcal{L}(n, \mathbb{R}^n, i\mathbb{R}^n) \to \Omega\mathcal{L}(n+2, \mathbb{R}^{n+2}, i\mathbb{R}^{n+2})$$

given by direct sum with γ. If we do this k times, we can homotope the map $(\oplus\gamma)^{\circ k} \circ f'$ to factor through $\Omega_m(n+2k)$.

Going back with the homeomorphism to $\Omega\mathcal{L}(n)$ we see that the stabilization we did corresponds to having stabilized with

$$e^{i\pi t/2}\gamma(t) = e^{i\pi t}\mathbb{R} \oplus \mathbb{R}$$

k times. We have argued that we can homotope the map after such a stabilization to the following subspace

$$e^{i\pi t/2}\Omega_m = \{\gamma \mid \gamma(t) = e^{i\pi t}W \oplus W^\perp\},$$

so by further stabilizing with

$$\gamma_2(t) = (e^{-i\pi t}\mathbb{R})^{\oplus k},$$

one has in total stabilized f with something homotopic to a standard stabilization, because it is easy to get the two "twistings" to cancel out. We have now homotoped the map f, stabilized in the standard way, to a map into $\Omega_s(\mathcal{L}(n+3k))$.

The last statement in the lemma follows from the fact that these highly connected inclusions of Grassmannians into $\Omega\mathcal{L} \simeq \mathbb{Z} \times B\mathrm{O}$ used in the proof, are the standard way of identifying the stable bundle with the difference of two actual bundles. □

What we in fact proved was that the map is homotopic to a map factoring through $\Omega_s\mathcal{L}(n_2)$, where V_- is the trivial bundle of dimension k. This is equivalent to the fact that any stable bundle over a compact space can be written as the difference between a bundle and a trivial bundle.

If we want to use this together with lemmas 6.5 and 6.6, we need to calculate negative bundles for the quadratic form B_r^D whenever $S_\mathcal{L}$ is of product type and on standard form on the factor ΛD^{2k}.

Lemma 7.3. *Assume that r is odd and large enough. For any point γ in $X = \mathcal{L}(n_1) \times \Omega_s\mathcal{L}(n_2) \subset \Lambda\mathcal{L}(n_1+n_2)$ we have an associated real quadratic form $B_r^\gamma \colon \mathbb{C}^{(n_1+n_2)r} \to \mathbb{R}$ defined by the Hessian of the finite reduction for $H_D = 0$ described in lemma 6.6. The negative eigenbundle over X of this quadratic form is isomorphic to*

$$\mathbb{R}^{(n_1+n_2)(r-2)+n_1} \oplus V_+ \oplus V_-^\perp,$$

where $V_-^\perp = V_+ \oplus V_0$.

We use the notation $V_-^\perp$ to emphasize that it is a complement bundle to V_-, making the bundle in the lemma isomorphic to $V_+ - V_-$ plus a trivial bundle of dimension $(n_1+n_2)(r-1)$.

Note that even though these quadratic forms have kernels, there is indeed a negative eigenbundle, because the kernel is the same for all the forms and thus defines a trivial bundle.

Proof: Given any $\gamma \in X$ we will describe a choice of subspace on which B_r^γ is negative, and argue that it has maximal dimension. It will be obvious that this choice is continuously dependent on γ, and we will argue that as a bundle over X it is indeed isomorphic to the one claimed in the lemma.

Because $\gamma(t) = L \oplus \gamma_1(t)$ with $L \subset \mathbb{C}^{n_1}$ and $\gamma_1(t) \subset \mathbb{C}^{n_2}$ we see that B_r^γ splits as a sum of $B_r^L \colon \mathbb{C}^{n_1 r} \to \mathbb{R}$ and $B_r^{\gamma_1} \colon \mathbb{C}^{n_2 r} \to \mathbb{R}$.

For now we restrict our attention to B_r^L. Note that for any quadratic form $x^T A x$, the set of critical points is precisely the kernel of A. We wish to define real vector spaces E_-, E_0 and E_+ on which B_r^L is negative, zero (to the second order) and positive respectively. Because $E_- \oplus E_0 \oplus E_+$ will be all of $\mathbb{C}^{n_1 r}$ we conclude that $\dim(E_-)$ is maximal.

Let $\rho = e^{i2\pi/r}$ be the standard r'th root of unity. Use this to define the real vector spaces E_m by

$$E_m = \{(b\rho^{mj})_{j\in\mathbb{Z}_r} \mid b \in \mathbb{C}^{n_1}\} \subset \mathbb{C}^{n_1 r},$$

for any $m \in \mathbb{Z}_r$. The action of $\mathrm{U}(n_1)$ preserves E_m and $\oplus_m E_m = \mathbb{C}^{n_1 r}$. It is convinient to think of these spaces as the finite version of the splitting of $L^2(S^1, \mathbb{C}^{n_1})$ into subspaces $E_m^{L^2} = \{be^{m2\pi it} \mid m \in \mathbb{Z}, t \in \mathbb{R}/\mathbb{Z}, b \in \mathbb{C}^{n_1}\}$.

Let M_- be the subset of $\mathbb{Z}_r$ containing the classes $[1], [2], \dots, [(r-1)/2]$, and $M_+ = -M_-$. These are disjoint and $\mathbb{Z}_r = M_- \cup M_+ \cup \{0\}$. Use this to define

$$E_- = \bigoplus_{m\in M_-} E_m$$
$$E_+ = \bigoplus_{m\in M_+} E_m.$$

We have already defined E_0 as the constant curves, which we know to be critical points of B_r^L, because for r large enough the only critical points of B_r^L are the 1-periodic orbits. One may extend the analogy to the splitting of $L^2(S^1, \mathbb{C}^{n_1}) \cong \Lambda\mathbb{C}^{n_1}$, by considering the actual action integral on this space for $H = 0$. Indeed, the action integral splits orthogonally on the splitting $E_m^{L^2}$ into eigen spaces with eigen values $-2\pi m$. So this finite version can be thought of as analagous to the spaces $E_m^{L^2}$ for $-(r-1)/2 \leq m \leq (r-1)/2$, and indeed the rest of the proof will reflect this analogy.

Since $H = 0$ we get no flow and the first term in the definition of the finite dimensional reduction vanishes (see the definition of the finite approximations in section 5). So B_r^L is in fact just a calculation of the symplectic area for the concatenation of the L-curves connecting z_j to z_{j+1} defined by L. The action of $\mathrm{U}(n_1)$ just rotates the curves and preserves the symplectic area and the spaces E_m, so we can assume that $L = i\mathbb{R}^{n_1}$. This means that B_r^L splits as a sum $A_1 + A_2 \cdots + A_{n_1}$, where A_k only depends on the k'th complex coordinate of the z_j's. We now look at one A_k at a time. In fact we assume for the time being that $n_1 = 1$, so that we do not have to redefine E_m. This means that symplectic volume is now minus the normal area in $\mathbb{C}$.

We will need the following facts about the spaces E_m, which because of the assumption $n_1 = 1$ are subspaces of $\mathbb{C}^r$. For $(z_j)_{j\in\mathbb{Z}_r} \in E_m$ and $(w_j)_{j\in\mathbb{Z}_r} \in E_{m'}$ we have

$$\begin{aligned} (z_{j+1})_{j\in\mathbb{Z}_r} &\in E_m \\ (\overline{z_j})_{j\in\mathbb{Z}_r} &\in E_{-m} \\ \mathrm{Re}(z_j)_{j\in\mathbb{Z}_r} &\in E_m \oplus E_{-m} \\ \mathrm{Im}(z_j)_{j\in\mathbb{Z}_r} &\in E_m \oplus E_{-m} \\ (z_j \cdot w_j)_{j\in\mathbb{Z}_r} &\in E_{m+m'}. \end{aligned}$$

and if $m \neq 0$

$$\sum_j z_j = 0.$$

Notice that the second to last fact makes sense only because we have $n_1 = 1$.

Any vector $\overrightarrow{z} = (z_j)_{j\in\mathbb{Z}_r} \in E_-$ can be written as

$$z_j = \sum_{m\in M_-} \alpha_m \rho^{mj}$$

with $\alpha_m \in \mathbb{C}$. Since $L = i\mathbb{R}$, the L-curves are parallel to the real axis on the first geodesic piece and parallel to the imaginary axis on the second. So we have

$$B_r^L(\overrightarrow{z}) = \sum_j y_j(x_{j+1} - x_j),$$

where $z_j = x_j + iy_j$. Rewriting this we get:

$$\begin{aligned} 4B_r^L(\overrightarrow{z}) &= 2\sum_j((y_{j+1}+y_j)(x_{j+1}-x_j) - (y_{j+1}-y_j)(x_{j+1}-x_j)) \\ &= \sum_j(2(y_{j+1}+y_j)(x_{j+1}-x_j) - \mathrm{Im}((z_{j+1}-z_j)^2)) \qquad (19) \\ &= \sum_j 2(y_{j+1}+y_j)(x_{j+1}-x_j). \end{aligned}$$

The last equality holds because of the facts stated for the spaces E_m and because we restricted $\overrightarrow{z}$ to E_-, so that no term involving E_0 occurs in the expression.

This splitting of the sum into two terms has a geometric interpretation: Take the area enclosed by the curve defined by just connecting the points z_j by straight lines, but then subtract the area defined by the triangles that are defined by the L-curves and this straight line. So what we just argued was that the areas of the triangles cancel each other out (on E_-). Using our expression for $\overrightarrow{z}$ we also see that all terms involving products of terms with different values of m cancel out. This actually means that B_r^L *restricted to* E_- splits orthogonally on the subspaces E_m, $m \in M_-$. All we need now is to calculate B_r^L on each of the E_m's, but here the geometrical interpretation tells us that the area is simply

$$2B_r^L((\alpha_m \rho^{mj})_{j\in\mathbb{Z}_r}) = -\sum_j \|\alpha_m\|^2 \sin(2\pi m/r) = -r\|\alpha_m\|^2 \sin(2\pi m/r),$$

which is negative because $m \in M_-$.

The same arguments show that B_r^L is positive on E_+, but be warned: The spaces E_m is not an orthogonal splitting of B_r^L. This is because E_- and E_+ are not orthogonal, since canceling the terms really used the assumptions on M_- and M_+.

If n_1 is not 1 it is easy to see that the E_m's are just direct sums of the E_m^i's and thus see that a E_- is a maximal negative subspace for B_r^L.

We now turn our attention to $A = B_r^{\gamma_1}$. It is easy to see that because of the form of γ_1, we can again split A into a sum of

$$\begin{aligned} A_+ &\colon (\mathbb{C}\otimes V_+)^r \to \mathbb{R} \\ A_0 &\colon (\mathbb{C}\otimes V_0)^r \to \mathbb{R} \\ A_- &\colon (\mathbb{C}\otimes V_-)^r \to \mathbb{R}, \end{aligned}$$

where $(\mathbb{C}\otimes V_+)\oplus(\mathbb{C}\otimes V_0)\oplus(\mathbb{C}\otimes V_-) = \mathbb{C}^{n_2}$ since $V_+\oplus V_0 \oplus V_- = \mathbb{R}^{n_2}$.

In the case of A_0 we see that this is very similar to the previous case of B_r^L. In fact, using an isometry $V_0 \cong \mathbb{R}^k$, we see that A_0 is actually the same as B_r^L. So a choice of maximal negative subspace for A_0 is

$$\begin{aligned} E_-^0 &= \bigoplus_{m\in M_-} E_m^0 \\ E_m^0 &= \{(b\rho^{mj})_{j\in\mathbb{Z}_r} \mid b \in \mathbb{C}\otimes V_0\}. \end{aligned}$$

Similarly, we take an isometry $V_+ \simeq \mathbb{R}^n$. We now split this as a sum and assume again that $n = 1$, and find a maximal negative subspace.

Recall the geometric interpretation of splitting B_r^L into two terms in equation (19). We see that the difference between having L constantly equal to $i\mathbb{R}$ and having the Lagrangian curve $t \mapsto e^{i\pi t}$, is only that the triangles, whose areas we have to subtract, are different. The area of the triangle defined by z_{j+1} and z_j and some Lagrangian L is given by

$$4\,\mathrm{area} = \mathrm{Im}(u^2(z_{j+1}-z_j)^2),$$

where u^{-1} is a unit vector in $\mathbb{C}$ with real span L. This means that the expression for A_+ is

$$4A_+(\overrightarrow{z}) = \sum_j (2(y_{j+1}+y_j)(x_{j+1}-x_j) - \mathrm{Im}(\rho^{-j}(z_{j+1}-z_j)^2)),$$

and this time the last sum does not sum to zero, and there is no symmetry between the previously defined positive and negative spaces. This is because multiplication by ρ^{-j} takes us from E_m to E_{m-1}.

We first prove that the spaces E_m are in fact orthogonal with respect to the bilinear form defined by the first part of the sum - that is, define:

$$\begin{aligned} Q(\overrightarrow{z}) &= \sum_j (y_{j+1}+y_j)(x_{j+1}-x_j) \\ q(\overrightarrow{z_1},\overrightarrow{z_2}) &= Q(\overrightarrow{z_1}+\overrightarrow{z_2}) - Q(\overrightarrow{z_1}) - Q(\overrightarrow{z_2}). \end{aligned}$$

By once again using the properties of the spaces E_m, we immediately see that E_m is orthogonal to $E_{m'}$ if $m \neq \pm m'$, so all we need to consider is $\overrightarrow{z_1} = \alpha_m \rho^{mj}$ and $\overrightarrow{z_2} = \alpha_{-m}\rho^{-mj}$ and prove that $q(\overrightarrow{z_1},\overrightarrow{z_2}) = 0$. By expanding the expression for $Q(\overrightarrow{z_1}+\overrightarrow{z_2})$ and comparing with $Q(\overrightarrow{z_2})+Q(\overrightarrow{z_1})$, we are left with two terms that are symmetric in changing the sign of m. Call these T_1 and T_2. We reduce one of these terms by

$$\begin{aligned} T_1 = \frac{1}{4i}\sum_j \Big(\alpha_m\rho^{mj}(\rho^m+1) - \overline{\alpha_m}\rho^{-mj}(\rho^{-m}+1)\Big)\cdot \\ \Big(\alpha_{-m}\rho^{-mj}(\rho^{-m}-1) + \overline{\alpha_{-m}}\rho^{mj}(\rho^m-1)\Big), \end{aligned}$$

and by using the properties we get

$$\begin{aligned} T_1 &= \frac{r}{4i}\Big(\alpha_m\alpha_{-m}(\rho^m+1)(\rho^{-m}-1) - \overline{\alpha_m\alpha_{-m}}(\rho^{-m}+1)(\rho^m-1)\Big) \\ &= \frac{r}{2}\operatorname{Im}\big(\alpha_m\alpha_{-m}(\rho^m+1)(\rho^{-m}-1)\big) = \frac{r}{2}\operatorname{Im}\big(\alpha_m\alpha_{-m}(\rho^{-m}-\rho^m)\big). \end{aligned}$$

Since T_2 was the same but with $-m$ and m interchanged we see that

$$T_1 + T_2 = 0.$$

Now we look at what happens on the previously positive subspace - that is, assume that $\overrightarrow{z} \in E_+$. The first part of the sum is the same as before. Looking at the last part we see that when summing we only get a contribution from the E_0 part of $\rho^{-j}(z_{j+1}-z_j)^2$. This means that we only get a contribution from the $E_1 = E_{-(r-1)}$ part of $(z_{j+1}-z_j)^2$. Since $(z_{j+1}-z_j)$ is in E_+, we only get a contribution from the $E_{-(r-1)/2}$ part of it. Define $m_1 = [(r+1)/2] = [-(r-1)/2] \in \mathbb{Z}_r$. From all this we get

$$2A_+(\overrightarrow{z}) = \Big(-\sum_{m\in M_+} r\|\alpha_m\|^2\sin(2\pi m/r)\Big) - r\operatorname{Im}\Big((\alpha_{m_1}(\rho^{m_1}-1))^2\Big)$$

for $\overrightarrow{z} \in E_+$. For r large enough we will have ρ^{m_1} approximately -1, so we get

$$2A_+(\overrightarrow{z}) \approx \Big(-\sum_{m\in M_+} r\|\alpha_m\|^2\sin(2\pi m/r)\Big) - 4r\operatorname{Im}\Big((\alpha_{m_1})^2\Big).$$

Denote by $Q \subset E_+$ the one-dimensional subspace where $\operatorname{Im}(\alpha_{m_1}^2) = \|\alpha_{m_1}\|^2$ and $\alpha_m = 0$, $m \neq m_1$. Also denote by $Q' \subset E_+$ the $(r-2)$-dimensional subspace given by $\operatorname{Im}(\alpha_{m_1}^2) = -\|\alpha_{m_1}\|^2$. We now have $Q \oplus Q' = E_+$, and because the positive term coming from the sum is much smaller numerically than the "new" term on Q, we see that A_+ is negative on Q and positive on Q'.

The kernel of A_+ is the set of constant curves because it is still a finite approximation for $H_D = 0$.

Now assume that $\overrightarrow{z}$ is in the space $E_- \oplus Q$. Because the E_m's were orthogonal with respect to the first part of the expression we get the same first term. We also notice that the second part of the sum is again non-vanishing only when dealing with products from $E_{-(r-1)/2}$, so in fact we get a very similar expression:

$$2A_+(\overrightarrow{z}) \approx \Big(- \sum_{m \in M_- \cup \{m_1\}} r\|\alpha_m\|^2 \sin(2\pi m/r)\Big) - 4r \operatorname{Im}\Big((\alpha_{m_1})^2\Big).$$

Again, because of the assumption that $\|\alpha_{m_1}\|^2 = \operatorname{Im}(\alpha_{m_1}^2)$, we see that this is negative. There is one positive term in the sum, but this as before is smaller numerically than the "new" negative term.

This leads us to define:

$$\begin{aligned} E_-^+ &= \Big(\bigoplus_{m \in M_-} E_m^+\Big) \oplus Q^+ \\ E_m^+ &= \{(b\rho^{mj})_{j \in \mathbb{Z}_r} \mid b \in \mathbb{C} \otimes V_+\} \\ Q^+ &= \{(b\rho^{m_1 j})_{j \in \mathbb{Z}_r} \mid b \in \mathbb{C} \otimes V_+, \|b\|^2 = \operatorname{Im}(b^2)\}. \end{aligned}$$

From the above arguments it is clear that E_-^+ is a maximal negative subspace for A_+. Note that Q^+ is canonically isomorphic to V_+.

We also define:

$$\begin{aligned} E_-^- &= \Big(\bigoplus_{m \in M_- - \{m_1\}} E_m^-\Big) \oplus Q^- \\ E_m^- &= \{(b\rho^{mj})_{j \in \mathbb{Z}_r} \mid b \in \mathbb{C} \otimes V_-\} \\ Q^- &= \{(b\rho^{(m_1-1)j})_{j \in \mathbb{Z}_r} \mid b \in \mathbb{C} \otimes V_-, \|b\|^2 = -\operatorname{Im}(b^2)\}. \end{aligned}$$

A similar calculation shows us that E_-^- is a maximal negative subspace for A_-. Note that as before Q^- is isomorphic to V_-, but in this case it is more important that $E_-^- \oplus V_-$ is canonically isomorphic to

$$\bigoplus_{m \in M_-} E_m^-,$$

which leads us to introduce the notation $E_-^- \cong (\oplus_{m \in M_-} E_m^-) \ominus V_-$.

Putting all this together we see that

$$\begin{aligned} W &= E_- \oplus E_-^+ \oplus E_-^0 \oplus E_-^- \cong \Big(\bigoplus_{m \in M_-} W_m\Big) \oplus V_+ \ominus V_- \\ W_m &= \{b\rho^{mj} \mid b \in \mathbb{C}^{n_1+n_2}\} \end{aligned}$$

is a choice of maximal negative subspace. □

8 The Viterbo Transfer as a Map of Spectra

Let L and N be closed smooth n-manifolds and let $j\colon L \to T^*N$ be an exact Lagrange embedding, i.e. the canonical one form λ_N on T^*N is pulled back to and exact 1-form by j. In this section we construct the Viterbo transfer (see [10]) as a map of spectra

$$(\Lambda j)_!\colon \Lambda N^{-TN} \to \Lambda L^{-TL+\eta},$$

where η is the Maslov bundle defined in definition 7.1, and ΛN^{-TN} is notation for the Thom-spectrum defined by a complement bundle to $TN = \mathrm{Ev}_0^* TN$. The construction is similar to that of Viterbo, but the lemmas in the previous sections make it much easier to control and understand the actual Thom spaces, and thus get the map as a spectrum map.

Start by giving both N and L Riemannian metrics. The Darboux-Weinstein Theorem tells us (see e.g. [7]) that for a small $\varepsilon_1 > 0$, we can symplectically extend j to a map $j\colon T^*_{2\varepsilon_1}L \to T^*N$. To distinguish between coordinates in T^*N and T^*L, we denote them by (q_N, p_N) and (q_L, p_L) respectively. It is very important for the construction that exactness of the embedding implies that $p_N dq_N - p_L dq_L = \lambda_N - \lambda_L$ defined on $T^*_{2\varepsilon_1}L$ is exact. It is closed but not exact for a non-exact Lagrange embedding. This implies that the two action integrals $\int_\gamma \lambda_N - Hdt$ and $\int_\gamma \lambda_L - Hdt$ are equal on closed curves in $T^*_{2\varepsilon_1}L$. This means that if we have a Hamiltonian on T^*N, which restricted to our neighborhood of L depends only on $\|p_L\|$, then we can use the method in remark 3.1 to calculate the action integral on closed 1-periodic orbits. In the following definition it is important to keep this method in mind.

First we define H near L. Here we define the Hamiltonian in terms of the coordinates (q_L, p_L). In fact on the set $T^*_{\varepsilon_1}L$ we want it to as in remark 5.1. So we want it to be a function of $\|p_L\|$, convex and linear with slope μ_L outside some small δ-neighborhood of the zero section. As before, we assume that μ_L is not the length of any geodesic in L.. On the rest of $T^*_{2\varepsilon_1}L$ we want it to be concave and "level off" to be a constant c near the boundary (see figure 6). We want this leveling off to happen so fast that any 1-periodic orbit in $T^*_{2\varepsilon_1}L - T^*_{\varepsilon_1}L$ has action below $-\delta\mu_L$. By using lemma 4.4, this can be done if δ is small enough, as seen in figure 6, where $\mu'_L < \mu_L$ is the maximum of



Figure 6: Definition of H.

geodesic lengths less than μ_L. It is clear that the larger μ_L is, the larger c becomes.

Outside $T^*_{2\varepsilon_1}L$, we define H to be the constant c on $T^*_R N - T^*_{2\varepsilon_1}L$ for a fixed $R > 0$, such that the embedding of $T^*_{2\varepsilon_1}L$ is inside $T^*_R N$. Outside $T^*_R N$, we define it to be a function of $\|p_N\|$, convex and linear outside some $R' > R$ with slope μ_N. Again we want the action of the 1-periodic orbits coming from this part to be less than $-\delta\mu_L$. So the larger c is, the larger we can choose μ_N (see figure 6).

We want to pick all the parameters such that we can use the results in all of the preceding sections, but first we concentrate on section 4, and assume that T^*N has the induced metric. Notice that $C_1 = \max(k\mu_L, \mu_N)$ works as the bound on the gradient of H. The k is needed because we used the induced metric from L to define H near L and this bound is in the metric induced from N. Similarly we can for small δ assume that δ^{-1} times some constant is the upper bound C_2 on the covariant Hessian of H. For small δ we can further assume that $C_2 > C_1$. So using proposition 4.7, we conclude that there exist a $K > 0$ and $\delta_0 > 0$ such that: If $0 < \delta < \delta_0$ and $r > K\delta^{-1}$ then

$$I(A_r, X) = \mathrm{Th}(T\Lambda_r^{\mu_N} N),$$

where X is the pseudo-gradient defined in the section.

Because of the way we defined H, we see that the natural quotient map, defined in the end of section 2, gives a map from the total index to the index $I_\delta = I^{+\delta\mu_L}_{-\delta\mu_L}(A_r, X)$. Recall that the action of H restricted to $T^*_{\varepsilon_1}L$ has all of its critical values in this interval, and all other critical values are below $-\delta\mu_L$.

Proposition 5.9 tells us that by possibly making K larger and choosing a β and then making δ_0 smaller, we get a good index pair for the index I_δ on the subset $\Lambda_{r,\beta}T^*_{\varepsilon_1}L$, provided $r \in [K\delta^{-1}, 2K\delta^{-1}]$ and $0 < \delta < \delta_0$. For this to make sense, we recall that X in section 4 was defined to be the gradient of A_r when $\max_j \|\varepsilon_{q_j}\|$ were small, so we may assume that X is the gradient of A_r on the subset $\Lambda_{r,\beta}T^*_{\varepsilon_1}L$, which was the assumption in section 5. Furthermore, a good pair defined inside a subset is also a good pair on the entire set, provided that it contains all the critical points, and we already accounted for that.

We have two metrics on $T^*_{\varepsilon_1}L$: One from T^*N, which we will denote g_N, and one induced from L, which we will denote g_L. We also have two canonical sections in the bundle in equation (10) (with N replaced by L), which come from taking the vertical directions with respect to T^*N and T^*L. We denote these $S^N_{\mathcal{L}}$ and $S^L_{\mathcal{L}}$ respectively. This data produces two finite reductions

$$A^1_r = A_r^{g_N, S^N_{\mathcal{L}}} : \Lambda^{g_N}_{r,\beta}T^*_{\varepsilon_1}L \to \mathbb{R}$$
$$A^2_r = A_r^{g_L, S^L_{\mathcal{L}}} : \Lambda^{g_L}_{r,\beta}T^*_{\varepsilon_1}L \to \mathbb{R}.$$

The next proposition relates the two homotopy indices.

Proposition 8.1. *Assume that r is odd and large enough. The total index of A^1_r, which we denoted I_δ, is stably equivalent to*

$$\mathrm{Th}((T\Lambda_r^{\mu_L} L) \oplus \eta),$$

where η is the Maslov bundle defined in definition 7.1.

Proof: First we argue that the homotopy index does not depend on the metric: use the convex change to get a smooth family g_t of metrics relating one with the other. Make sure that $K > 0$ and $\beta > 0$ are large enough for proposition 5.9 to work for all metrics g_t. Then fix δ_0 such that the lemma works for this K, β and all metrics g_t. We now have good pairs for all t, but it is a problem that the spaces $\Lambda^{g_t}_{r,\beta} T^*_{\varepsilon_1} L$ depend on t. However, this can be resolved locally: Given any $t \in [0,1]$ we want to argue that the homotopy index is constant in a small neighborhood of t even though the manifold changes. This is because the boundary of the manifold is given by a continuous function on a bigger manifold, so by going to a slightly smaller manifold still containing our good index pair, we can use lemma 2.4.

Let s_N and s_L be the sections in the bundle

$$\mathcal{L}(T(T^*_{\varepsilon_1} L)) \to T^*_{\varepsilon_1} L$$

given by the vertical foliations from of T^*N and T^*L respectively. Then $S^N_{\mathcal{L}} = \Lambda s_N$ and $S^L_{\mathcal{L}} = \Lambda s_L$.

Redefine A^1_r as the finite approximation defined by $S^N_{\mathcal{L}}$ but using the metric induced from L. As we just argued, this does not change the index.

We stabilize the section $S^N_{\mathcal{L}}$ to the section of product type (see definition 6.1)

$$S^N_{\mathcal{L},k} = S^N_{\mathcal{L}} \oplus i\mathbb{R}^k$$

and get a new finite dimensional approximation from section 6. Lemma 6.5 tells us that since we chose this section to be independent of the point in $\Lambda_{r,\beta} T^*_{\varepsilon_1} L$ on the second factor, this new approximation has as its total index the m-fold suspension of I_δ for some m.

If we chose k large enough we can as in the beginning of section 7 homotope the difference of s_N and s_L into $\mathcal{L}(k)$ - that is, we can homotope s_N through sections in $\mathcal{L}T(T^*_{\varepsilon_1} L \times D^{2k}) \to T^*_{\varepsilon_1} L \times D^{2k}$ to

$$s_L \oplus J_*,$$

where s_L and J_* do not depend on the point in D^{2k}. So when looping this we get a section of product type.

Because we can assume that this homotopy is smooth, by looping it we get a homotopy of sections in the bundle (10) with the bounds in equation (11). However, we need lemma 7.3 to identify the negative eigenbundle of the quadratic form given the finite approximations defined by stabilizing s_L with J_*. For this we need to homotope ΛJ_* to the subset of curves in lemma 7.3, and this can be done by lemma 7.2 if k was chosen large enough. The catch is that we need this homotopy to fulfill the bounds in equation (11). If we do not have these bounds, we cannot argue that the homotopy index is constant during the homotopy, because we do not know if good index pairs exist. The map, we are homotoping ΛJ_* to, is not the loop of a map. So we must argue in some other way that this can be done within the bounds. Indeed, there exists a deformation retraction $f_t, t \in [0,1]$ of the set of curves with energy less than β onto a finite dimensional compact manifold satisfying a bound like the second one in (11)[3]. The homotopy $f_t \circ \Lambda J_*$ fulfills the bounds, because ΛJ_*

---

[3] just pick an s such that $\beta/\sqrt{s}$ is smaller than the injective radius and contract onto piecewise geodesics

does and f_t fulfills the last of them, and maps the set with energy less than β to itself. The image of f_1 is a compact manifold, so homotoping ΛJ_* only on this subspace into the set we need it to lie in, will ensure the needed bounds for the composition.

This homotopy is a compact family of sections, so there is a $\delta_0 > 0$ small enough to make proposition 5.9 and lemma 6.2 work for all of them simultaneously. So we use lemma 6.5 to conclude that the index I_δ is the relative Thom suspension of the total index of A_r^2 by the negative eigenbundle of the quadratic form added. We then use lemma 6.6 and lemma 7.3 to conclude that this bundle is a trivial bundle of dimension m plus the Maslov bundle. □

Corollary 8.2. *The index quotient map to I_δ induces a map of spectra*

$$(\Lambda^{\mu_N} N)^{-TN} \to (\Lambda^{\mu_L} L)^{-TL \oplus \eta},$$

where $\Lambda^\mu X$ is the loop space of curves with lengths less than μ in the Riemannian metric on X.

Proof: First we note that the space

$$\mathrm{Th}(T\Lambda_r^{\mu_N} N)$$

can by parallel transport along the curve be identified with

$$(\Lambda_r^{\mu_N} N)^{\mathrm{Ev}_{z_0}^* TN^{\oplus r}}.$$

Here we switch notation for Thom bundles, because the name of the base space is not included in the bundle name. This identification makes sense in light of lemma 4.6.

Let $\psi \to A$ be any bundle over A where (A, B) is an index pair defining the total index of $A_r \colon T^*\Lambda_r N \to \mathbb{R}$. Then any quotient map $f \colon A/B \to A/B'$ has a relative Thom suspension using ψ defined by

$$D\psi/(D\psi_{|B} \cup S\psi) \to D\psi/(D\psi_{|B'} \cup S\psi).$$

If A/B is already a Thom space of some bundle $\psi' \to Y$, then the left hand side is the Thom space of $\psi' \oplus \psi_{|Y} \to Y$. Similarly for the right hand side. So using lemma 4.6 but adding on top of the index quotient a copy of a complement to the bundle $\mathrm{Ev}_{q_0}^* TN$, we do a standard suspension of the indices instead of the Thom suspension by the bundle. So the new quotient map obtained is just the suspension of the old one. This, however, looks a bit confusing considering the previous lemma: When increasing r by 2 we Thom suspended the index I_δ by the bundle $2\,\mathrm{Ev}_{q_0}^* TL$, but we just added two times a complement to $\mathrm{Ev}_{q_0}^* TN$, so why should this be a suspension? This can be explained by the fact that since $T_{\varepsilon_1}^* L$ is embedded into T^*N, they have isomorphic tangent bundles and two copies of TL is thus isomorphic to two copies of the pullback of TN.

Since the previous lemma worked only for r odd we see that removing two copies of $\mathrm{Ev}_{q_0}^* TN$ and two copies of $\mathrm{Ev}_{q_0}^* TL$ on each side of the map

$$\mathrm{Th}(T\Lambda_r^{\mu_N} N) \to \mathrm{Th}(T\Lambda_r^{\mu_L} L \oplus \eta)$$

cannot fully convert all the tangent spaces to suspensions on both sides simultaneously, so we choose the convention of removing an extra on each side, making both sides into spaces representing the wanted spectra. □

As noted earlier, the larger μ_L is, the larger we can choose μ_N, which is a good indication that these maps glue together into something non-trivial.

Theorem 1. *The Viterbo transfer can be realized as a map of spectra*

$$(\Lambda j)_!\colon (\Lambda N)^{-TN} \to (\Lambda L)^{-TL+\eta}$$

Proof: The map is the same as the Viterbo transfer because the construction is the same. The difference is that in our construction we actually calculate the homotopy type of I_δ (not the same name as in [10]) and add appropriate bundles to both sides.

We need to argue that the maps from the previous corollary are compatible with inclusions when increasing either μ_L or μ_N. Since we already proved that the quotients induce the maps, we only need to prove that they are compatible with inclusions.

First we establish two facts about the inclusion maps for Hamiltonians with two different slopes $\mu_1 < \mu_2$ not geodesics lengths. For the Hamiltonian H_1 in figure 7 we can use lemma 4.7, lemma 4.5 and the note right after the proof of lemma 4.5 to conclude that the inclusion of homotopy indices $I^{a}_{-1}(A_r, X)$ into the total index induces the Thom suspension of the inclusions defined by including curves with length less than μ_1 into the set of curves with length less than μ_2.



Figure 7: The Hamiltonian H_1.

For the approximation associated to the Hamiltonian H_2 in figure 8 we have a natural quotient map from the total index to $I^{+\mu\delta}_{-\mu\delta}(A_r, X)$. We will show that this is homotopic to the inclusion as above. In the case of H_2' we can, by making the part with slope μ_1 long enough, assume that the new bend has its associated critical points above $\mu_2\delta$. The inclusion of $I^{\mu_2\delta}_{a}(A_r, X)$ into the total index is the same as the previous inclusion. Now by shortening the linear part with slope μ_1, and then letting the bends cancel out, we remove



Figure 8: Hamiltonians H_2 and H_2'.

the critical points with critical value less than $-\mu_2\delta$. If we think of this in terms of Morse theory and CW-complexes, we see that the new cells glued on in the inclusion effectively kill of the subcomplex $I_a^{-\mu_2\delta}(A_r, X)$, so collapsing this subcomplex is homotopic to the inclusion. This argument can be made precise by doing a small perturbation of A_r to make it Morse. So the two maps are indeed homotopic.

Now look at the problem of increasing μ_L. This can be done by a homotopy that simply multiplies the part of H close to L by $a_t \in \mathbb{R}$ increasing in t where $a_0 = 1$ and a_1 is the ratio between the new and the old μ_L. Outside $T^*_{2\varepsilon_1}L$ we simply translate H upwards so we do not change the slope μ_N. During this process we move the critical points, and when $a_t\mu_L$ is a geodesic length we create new critical points. We create two critical points per geodesic with this length: One down by the bottom bend very close to the zero section of L, and one up in the concave part of H. At their creation they have the same critical value, which is larger than all the other critical values. We assume



Figure 9: Lowering μ_N.

that our bounds for the index pair are chosen such that this is within I_δ. One of these new critical values stays above zero, but the other (from the concave part) moves below zero and through $-\mu_L\delta$, effectively collapsing this part of the homotopy index, but as we saw, this collapse induces the inclusion.

The case of changing μ_N is much easier, because by looking at the Hamiltonian on figure 9, taking the quotient from the total index to I_δ is the composition of two quotients where one is the inclusion and the other is the quotient for the larger μ_N. $\square$

Bibliography

[1] Alberto Abbondandolo and Matthias Schwarz. On the Floer homology of cotangent bundles. *Comm. Pure Appl. Math.*, 59(2):254–316, 2006.

[2] Marc Chaperon. Une idée du type "géodésiques brisées" pour les systèmes hamiltoniens. *C. R. Acad. Sci. Paris Sér. I Math.*, 298(13):293–296, 1984.

[3] Ralph L. Cohen and John D. S. Jones. A homotopy theoretic realization of string topology. *Math. Ann.*, 324(4):773–798, 2002.

[4] Charles Conley. *Isolated invariant sets and the Morse index*, volume 38 of *CBMS Regional Conference Series in Mathematics*. American Mathematical Society, Providence, R.I., 1978.

[5] Kenji Fukaya, Paul Seidel, and Ivan Smith. Exact Lagrangian submanifolds in simply-connected cotangent bundles. *arXiv:math/0701783v1*, 2007.

[6] A. B. Givental′. Periodic mappings in symplectic topology. *Funktsional. Anal. i Prilozhen.*, 23(4):37–52, 96, 1989.

[7] Dusa McDuff and Dietmar Salamon. *Introduction to symplectic topology*. Oxford Mathematical Monographs. The Clarendon Press Oxford University Press, New York, second edition, 1998.

[8] J. Milnor. *Morse theory*. Based on lecture notes by M. Spivak and R. Wells. Annals of Mathematics Studies, No. 51. Princeton University Press, Princeton, N.J., 1963.

[9] David Nadler. Microlocal branes are constructible sheaves. *arXiv: math/0612399v2*, 2007.

[10] Claude Viterbo. Exact Lagrange submanifolds, periodic orbits and the cohomology of free loop spaces. *J. Differential Geom.*, 47(3):420–468, 1997.